

On the Decidability of Monadic Second-Order Logic with Arithmetic Predicates

Valérie Berthé
berthe@irif.fr
Université de Paris, IRIF, CNRS
Paris, France

Toghrol Karimov
Joris Nieuwveld
Joël Ouaknine
Mihir Vahanwala
toghs@mpi-sws.org
jnieuwve@mpi-sws.org
joel@mpi-sws.org
mvahanwa@mpi-sws.org
Max Planck Institute for
Software Systems
Saarbrücken, Germany

James Worrell
jbw@cs.ox.ac.uk
University of Oxford, Department of
Computer Science
Oxford, United Kingdom

ABSTRACT

We investigate the decidability of the monadic second-order (MSO) theory of the structure $\langle \mathbb{N}; <, P_1, \dots, P_k \rangle$, for various unary predicates $P_1, \dots, P_k \subseteq \mathbb{N}$. We focus in particular on ‘arithmetic’ predicates arising in the study of linear recurrence sequences, such as fixed-base powers $\text{Pow}_k = \{k^n : n \in \mathbb{N}\}$, k -th powers $N_k = \{n^k : n \in \mathbb{N}\}$, and the set of terms of the Fibonacci sequence $\text{Fib} = \{0, 1, 2, 3, 5, 8, 13, \dots\}$ (and similarly for other linear recurrence sequences having a single, non-repeated, dominant characteristic root). We obtain several new unconditional and conditional decidability results, a select sample of which are the following:

- The MSO theory of $\langle \mathbb{N}; <, \text{Pow}_2, \text{Fib} \rangle$ is decidable;
- The MSO theory of $\langle \mathbb{N}; <, \text{Pow}_2, \text{Pow}_3, \text{Pow}_6 \rangle$ is decidable;
- The MSO theory of $\langle \mathbb{N}; <, \text{Pow}_2, \text{Pow}_3, \text{Pow}_5 \rangle$ is decidable assuming Schanuel’s conjecture;
- The MSO theory of $\langle \mathbb{N}; <, \text{Pow}_4, N_2 \rangle$ is decidable;
- The MSO theory of $\langle \mathbb{N}; <, \text{Pow}_2, N_2 \rangle$ is Turing-equivalent to the MSO theory of $\langle \mathbb{N}; <, S \rangle$, where S is the predicate corresponding to the binary expansion of $\sqrt{2}$. (As the binary expansion of $\sqrt{2}$ is widely believed to be normal, the corresponding MSO theory is in turn expected to be decidable.)

These results are obtained by exploiting and combining techniques from dynamical systems, number theory, and automata theory.

CCS CONCEPTS

• Theory of computation \rightarrow Logic and verification.

KEYWORDS

Monadic second-order logic, linear recurrence sequences, toric words, cutting sequences, decidability

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

Conference’17, July 2017, Washington, DC, USA

© 2024 Association for Computing Machinery.

ACM ISBN 978-x-xxxx-xxxx-x/YY/MM... \$15.00

<https://doi.org/10.1145/nnnnnnn.nnnnnnn>

ACM Reference Format:

Valérie Berthé, Toghrol Karimov, Joris Nieuwveld, Joël Ouaknine, Mihir Vahanwala, and James Worrell. 2024. On the Decidability of Monadic Second-Order Logic with Arithmetic Predicates. In *Proceedings of ACM Conference (Conference’17)*. ACM, New York, NY, USA, 17 pages. <https://doi.org/10.1145/nnnnnnn.nnnnnnn>

1 INTRODUCTION

Büchi’s seminal 1962 paper [9] established the decidability of the monadic second-order (MSO) theory of the structure $\langle \mathbb{N}; < \rangle$, and in so doing brought to light the profound connections between mathematical logic and automata theory. Over the ensuing decades, considerable work has been devoted to the question of which expansions of $\langle \mathbb{N}; < \rangle$ retain MSO decidability. In other words, for which unary predicates P_1, \dots, P_k is the MSO theory of $\langle \mathbb{N}; <, P_1, \dots, P_k \rangle$ decidable?¹ Here by unary predicate we mean a fixed set of non-negative integers $P \subseteq \mathbb{N}$. Taking, for example, P to be the set of prime numbers, Büchi and Landweber [10] observed in 1969 that a proof of decidability of the MSO theory of $\langle \mathbb{N}; <, P \rangle$ would “seem very difficult”, as it would *inter alia* enable one (at least in principle) to settle the twin prime conjecture. (Decidability was subsequently established assuming Schinzel’s hypothesis H [5].)

The set of prime numbers is, of course, highly intricate. In 1966, Elgot and Rabin [15] considered a large class of simpler predicates of ‘arithmetic’ origin, such as, for any fixed k , the set $\text{Pow}_k = \{k^n : n \in \mathbb{N}\}$ of powers of k , and the set $N_k = \{n^k : n \in \mathbb{N}\}$ of k -th powers. For any such predicate P , they systematically established decidability of the MSO theory of $\langle \mathbb{N}; <, P \rangle$ by using (in modern parlance) a notion of *effective profinite ultimate periodicity* (essentially an automata-theoretic concept). Many years later, the theory was substantially developed and extended by Carton and Thomas [12], Rabinovich [28], and Rabinovich and Thomas [29], among others. A related key concept is that of *effective almost periodicity*, introduced in the 1980s by Semënov [32], and recently brought to bear in the MSO model checking of linear dynamical systems [20].

It is notable that whilst Elgot and Rabin establish separately the decidability of the MSO theories, for example, of $\langle \mathbb{N}; <, \text{Pow}_2 \rangle$ and

¹The restricted focus on *unary* (or *monadic*) predicates is justified by the fact that most natural non-unary predicates immediately lead to undecidability.

$\langle \mathbb{N}; <, \text{Pow}_3 \rangle$, they remain resolutely silent on the obvious joint expansion $\langle \mathbb{N}; <, \text{Pow}_2, \text{Pow}_3 \rangle$. This in hindsight is wholly unsurprising: there are various statements that one can express in the above theory whose truth values are highly non-trivial to determine: for example, for given fixed a, b , the assertion that there exist infinitely many powers of 3 whose distance to the next power of 2 is congruent to a modulo b . An immediate corollary of our first main result, Thm. 5.1, is that the MSO theory of $\langle \mathbb{N}; <, \text{Pow}_2, \text{Pow}_3 \rangle$ is indeed decidable. Although this is new, we should mention that decidability of the *first-order* theory of $\langle \mathbb{N}; <, \text{Pow}_2, \text{Pow}_3 \rangle$ has been known for over forty years, an important result of Semënov [31].

Looking over the last several decades' worth of research work on monadic second-order expansions of the structure $\langle \mathbb{N}; < \rangle$, it is fair to say that the bulk of the attention has focused on the addition of a single predicate P . The obvious reason is that whilst, in general, the decidability of single-predicate expansions of $\langle \mathbb{N}; < \rangle$ can usually be handled with automata-theoretic techniques alone, by reasoning about individual patterns in isolation, this is not the case when multiple predicates are at play simultaneously. Such collections of predicates can exhibit highly complex interaction patterns, which existing approaches are ill-equipped to handle.

In this paper, we show that key aspects of such interactions can be modelled in the theory of *dynamical systems*, and in particular via the notion of *toric words* [7]. In addition, we make use of number-theoretic tools to ensure effectiveness at various junctures of our algorithms. Some of our results are conditional: whereas Baker's theorem on linear forms in logarithms (alongside other tools) underpins the decidability of the MSO theory of $\langle \mathbb{N}; <, \text{Pow}_2, \text{Pow}_3 \rangle$, we are only able to show decidability of the MSO theory of $\langle \mathbb{N}; <, \text{Pow}_2, \text{Pow}_3, \text{Pow}_5 \rangle$ subject to Schanuel's conjecture, a central hypothesis in transcendental number theory. Intuitively, the reason is that whilst Baker's theorem suffices to handle interaction patterns of powers of 2 and powers of 3, the injection of powers of 5 into the mix exceeds the limits of contemporary number-theoretic knowledge.

Our paper contains two main results. Theorem 5.1 considers predicates arising from the sets of values achieved by certain *linear recurrence sequences*, generalising the predicates considered above. A simplified version of that result is as follows:

THEOREM 1.1. *Let $\rho_1, \dots, \rho_d > 1$ be natural numbers.*

- (1) *The MSO theory of $\langle \mathbb{N}; <, \text{Pow}_{\rho_1}, \dots, \text{Pow}_{\rho_d} \rangle$ is decidable assuming Schanuel's conjecture.*
- (2) *If $1/\log(\rho_1), \dots, 1/\log(\rho_d)$ are linearly independent over \mathbb{Q} , then the decidability is unconditional.*
- (3) *If each triple of distinct ρ_i, ρ_j, ρ_k is multiplicatively dependent, then the decidability is unconditional.*

Item (3) captures, for example, the decidability of the MSO theory of $\langle \mathbb{N}; <, \text{Pow}_2, \text{Pow}_3, \text{Pow}_6 \rangle$. Our second main result is Thm. 7.2, restated here:

THEOREM 1.2. *Let p, q, b, d be natural numbers such that $\eta = \sqrt[d]{p/q}$ is irrational, $P_1 = \{qn^d : n \in \mathbb{N}\}$, and $P_2 = \{pb^{nd} : n \in \mathbb{N}\}$. The decidability of the MSO theory of $\langle \mathbb{N}; <, P_1, P_2 \rangle$ is Turing-equivalent to that of the MSO theory of the base- b expansion of η .²*

²The representation of η in the base- b number system is an infinite word α over the finite alphabet $\Sigma = \{0, 1, \dots, b-1\}$ of digits. By MSO theory of an infinite word,

The underlying dynamical system here is of a symbolic nature: it consists of the base- b expansion of the irrational number η , which is a d -th root of a rational number. Such expansions are widely conjectured to be *normal*, and *a fortiori weakly normal*: every finite pattern of digits should occur infinitely often. As the MSO theory of any weakly normal word is decidable (Thm. 4.18), we obtain a conditional decidability result.

Note that when η is rational, we obtain unconditional decidability (Thm. 7.1), thanks to a composition result (Thm. 4.11) which we believe may be of independent interest. Here we state a simple corollary of Thm. 7.1:

COROLLARY 1.3. *For any positive integers b and d , the MSO theory of $\langle \mathbb{N}; <, \text{Pow}_{b^d}, N_d \rangle$ is decidable.*

For example, we recover from the above the decidability of the MSO theory of $\langle \mathbb{N}; <, \text{Pow}_4, N_2 \rangle$, mentioned in the abstract.

2 PRELIMINARIES

2.1 Words and Automata

By an alphabet Σ we mean a finite set of letters. For a finite or infinite word α and $n \in \mathbb{N}$, we write $\alpha(n)$ for the n th letter of α . Thus $\alpha = \alpha(0)\alpha(1)\dots$. We define $\alpha[n, m] := \alpha(n)\dots\alpha(m-1)$, and assuming α is infinite, $\alpha[n, \infty] := \alpha(n)\alpha(n+1)\dots$. We denote the length of a finite word w by $|w|$. A finite word $w \in \Sigma^*$ occurs at a position n in α if $\alpha[n, n+|w|] = w$. Such w is called a *factor* of α . We will often factorise $\alpha \in \Sigma^\omega$ as $\alpha = u_0u_1\dots$, where $u_i \in \Sigma^*$ for all $i \in \mathbb{N}$. Such a factorisation is uniquely determined by an increasing sequence $\langle k_n \rangle_{n=0}^\infty$ over \mathbb{N} such that $u_i = \alpha[k_i, k_{i+1}]$ for all i . Finally, consider $\alpha_i \in \Sigma_i^\omega$ for $1 \leq i \leq d$. The *product word* $\alpha := \alpha_1 \times \dots \times \alpha_d$ is defined by $\alpha(n) = (\alpha_1(n), \dots, \alpha_d(n)) \in \Sigma_1 \times \dots \times \Sigma_d$ for all n .

Let $\alpha \in \Sigma^\omega$. We say that α is

- (a) *effective* if for any $n \in \mathbb{N}$, $\alpha(n)$ can be effectively computed,
- (b) *weakly normal* if for every $w \in \Sigma^+$, w occurs as a factor of α infinitely often, and
- (c) *uniformly recurrent* if for every $w \in \Sigma^+$, either w does not occur in α , or there exists $R(w) \in \mathbb{N}$ such that w occurs in every factor of α of length $R(w)$. Equivalently, if w occurs in α , then there exists an integer $R(w)$ such that for all $N \in \mathbb{N}$, w occurs in $\alpha[N, N + R(w)]$.

Prominent examples of uniformly recurrent words include the Thue-Morse word [2, Chap. 1] and all Sturmian words [22, Chap. 2].

A *deterministic finite Muller automaton* \mathcal{A} over an alphabet Σ is given by a tuple $(Q, q_{\text{init}}, \delta, \mathcal{F})$, where Q is the (finite) set of states, q_{init} is the initial state, $\delta: Q \times \Sigma \rightarrow Q$ is the transition function, and \mathcal{F} is the acceptance condition consisting of subsets of Q . We denote by $\mathcal{A}(\alpha)$ the *run* of \mathcal{A} on α , which is the set of states visited when \mathcal{A} reads α . A word $\alpha \in \Sigma^\omega$ is *accepted* by \mathcal{A} if the set S of states appearing infinitely often in $\mathcal{A}(\alpha)$ is present in \mathcal{F} .

A *deterministic finite transducer* \mathcal{B} over an input alphabet Σ and an output alphabet Γ is given by $(R, r_{\text{init}}, \sigma)$, where R is the (finite) set of states, r_{init} is the initial state, and $\sigma: R \times \Sigma \rightarrow R \times \Gamma^*$ is the transition function. At every step, \mathcal{B} reads a letter from the input alphabet Σ , transitions to the next state, and outputs a finite word over the output alphabet Γ . We define $\sigma_R: R \times \Sigma \rightarrow R$ to be the

we mean the MSO theory of the structure $\langle \mathbb{N}; <, P_0, \dots, P_{b-1} \rangle$, where $P_i = \{n \in \mathbb{N} : \alpha(n) = i\}$ for $0 \leq i \leq b-1$.

function that returns the next state, and $\sigma_{\Gamma^*} : R \times \Sigma \rightarrow \Gamma^*$ to be the function that returns the output word. We denote by $\mathcal{B}(\alpha)$ the (possibly finite) word over Γ output by \mathcal{B} upon reading $\alpha \in \Sigma^\omega$.

Let \mathcal{A} be a finite automaton as above. By a *journey* on \mathcal{A} we mean an element of $J := Q \times Q \times 2^Q$. A path $q_0 q_1 q_2 \cdots q_n \in Q^{n+1}$ makes the journey (q_0, q_n, V) where V is the set of states occurring in the proper suffix $q_1 q_2 \cdots q_n$. If $n \geq 1$, then $q_n \in V$ necessarily, but q_0 may not belong to V . The unique journey a word $w \in \Sigma^*$ makes starting in $q \in Q$, denoted by $\text{jour}(w, q)$, is the journey made by $q_0 \cdots q_{|w|}$ where $q_{i+1} = \delta(q_i, w(i))$ for $1 \leq i < |w|$. The empty word makes journeys of the form (q, q, \emptyset) . If v makes the journey (q_1, q_3, V_1) and w makes the journey (q_3, q_2, V_2) , then vw makes the journey $(q_1, q_2, V_1 \cup V_2)$.

Next, we define the equivalence relation $\sim_{\mathcal{A}}$ as follows. Two words $v, w \in \Sigma^*$ are equivalent, denoted $v \sim_{\mathcal{A}} w$, if and only if the sets of journeys they can undertake (starting from various states) are identical. The equivalence is moreover a congruence: if $v \sim_{\mathcal{A}} w$ and $x \sim_{\mathcal{A}} y$, then $vx \sim_{\mathcal{A}} wy$. Observe that $\sim_{\mathcal{A}}$ is not the classical congruence associated with the automaton \mathcal{A} . Our choice, however, will be more convenient for technical reasons.

Since there are only finitely many equivalence classes of $\sim_{\mathcal{A}}$, the quotient of Σ^* by $\sim_{\mathcal{A}}$ is a finite monoid M . We use h to denote the natural morphism from Σ^* into M . The morphism h maps each letter to its equivalence class modulo $\sim_{\mathcal{A}}$. We also extend the function jour to take inputs from $M \times Q$: For equivalence class $m = [w]$ and state q , we define $\text{jour}(m, q) = \text{jour}(w, q)$. Finally, we will need the following lemma, whose proof is immediate.

LEMMA 2.1. *Let \mathcal{A} be an automaton as above and $\alpha \in \Sigma^\omega$ with factorisation $\alpha = u_0 u_1 \cdots \in \Sigma^\omega$, where $u_n \in \Sigma^*$ for all n . Then the word $\mathcal{A}(\alpha)$ can be decomposed as the concatenation of journeys*

$$(q_0, q_1, V_0)(q_1, q_2, V_1)(q_2, q_3, V_2) \cdots$$

where $q_0 = q_{\text{init}}$, $\text{jour}(u_n, q_n) = (q_n, q_{n+1}, V_n)$ for all n , and for every $q \in Q$, the state q appears infinitely often in $\mathcal{A}(\alpha)$ if and only if $q \in V_n$ for infinitely many $n \in \mathbb{N}$.

2.2 Monadic Second-Order Logic

Monadic second-order logic (MSO) is an extension of first-order logic that allows quantification over subsets of the universe. Such subsets can be viewed as unary (that is, monadic) predicates. We will only be interpreting MSO formulas over expansions of the structure $\langle \mathbb{N}; < \rangle$. For a general perspective on MSO, see [8].

Let $\mathbb{S} := \langle \mathbb{N}; <, P_1, \dots, P_m \rangle$ be a structure where each $P_i \subseteq \mathbb{N}$ is a unary predicate. We associate a language $\mathcal{L}_{\mathbb{S}}$ of terms and formulas with \mathbb{S} as follows. The terms of $\mathcal{L}_{\mathbb{S}}$ are the countably many constant symbols $\{0, 1, 2, \dots\}$, lowercase variable symbols that stand for elements of \mathbb{N} , and uppercase variable symbols that denote subsets of \mathbb{N} . The formulas of $\mathcal{L}_{\mathbb{S}}$ are the well-formed statements constructed from the built-in equality ($=$) and membership (\in) symbols, logical connectives, quantification over elements of \mathbb{N} (written Qx for a quantifier Q), and quantification over subsets (written QX for a quantifier Q). The MSO theory of the structure \mathbb{S} is the set of all sentences belonging to $\mathcal{L}_{\mathbb{S}}$ that are true in \mathbb{S} . The MSO theory of \mathbb{S} is *decidable* if there exists an algorithm that, given a sentence $\varphi \in \mathcal{L}_{\mathbb{S}}$, decides if φ belongs to the MSO theory of \mathbb{S} .

As an example, consider $\mathbb{S} = \langle \mathbb{N}; <, P \rangle$ where P is the set of all primes. Let $s(\cdot)$ be the successor function defined by $s(x) = y$ if and only if

$$x < y \wedge \forall z. x < z \Rightarrow y \leq z.$$

That is, $s(x) = x + 1$. Further let

$$\begin{aligned} \varphi(X) &:= 1 \in X \wedge 0, 2 \notin X \wedge \forall x. x \in X \Leftrightarrow s(s(x)) \in X, \\ \psi &:= \exists X: \varphi(X) \wedge \forall y. \exists z > y: z \in X \wedge P(z). \end{aligned}$$

The formula φ defines the subset $\{n: n \equiv 1 \pmod{3}\}$ of \mathbb{N} , and ψ is the sentence “there are infinitely many primes congruent to 1 modulo 3”, which is the case. At the time of writing, it is not known whether the MSO theory of the structure \mathbb{S} above is decidable.

The *Acceptance Problem* for an infinite word α , denoted Acc_α , is to determine, given a deterministic Muller automaton \mathcal{A} , whether \mathcal{A} accepts α . Let $P_1, \dots, P_d \subseteq \mathbb{N}$ be predicates and $\Sigma = \{0, 1\}^d$.

Definition 2.2. *The characteristic word of (P_1, \dots, P_d) , written $\alpha := \text{Char}(P_1, \dots, P_d) \in \Sigma^\omega$, is defined by $\alpha(n) = (b_{n,1}, \dots, b_{n,d})$ where $b_{n,i} = 1$ if $n \in P_i$ and $b_{n,i} = 0$ otherwise.*

The following is a seminal result of Büchi, through which he showed decidability of the MSO theory of $\langle \mathbb{N}; < \rangle$.

THEOREM 2.3 ([33, THMS. 5.4 AND 5.9]). *The decision problem of the MSO theory of the structure $\langle \mathbb{N}; <, P_1, \dots, P_d \rangle$ is Turing-equivalent to Acc_α , where α is the characteristic word of (P_1, \dots, P_d) .*

2.3 Algebraic Numbers

A complex number λ is algebraic if there exists $p \in \mathbb{Q}[x]$ such that $p(\lambda) = 0$. The set of algebraic numbers is denoted by $\overline{\mathbb{Q}}$. The unique irreducible monic polynomial that has λ as a root is called the *minimal polynomial* of λ . A *canonical representation* of an algebraic number λ consists of its minimal polynomial p and sufficiently accurate rational approximations of the real and imaginary parts of λ to distinguish it from the other roots of p . All arithmetic operations can be performed effectively on canonical representations of algebraic numbers [13, Sec. 4.2].

By a *multiplicative relation* of $(\lambda_1, \dots, \lambda_d) \in \mathbb{C}^d$ we mean $\mathbf{k} = (k_1, \dots, k_d) \in \mathbb{Z}^d$ such that $\lambda_1^{k_1} \cdots \lambda_d^{k_d} = 1$. We write

$$G := G_M(\lambda_1, \dots, \lambda_d)$$

for the set of all multiplicative relations of $(\lambda_1, \dots, \lambda_d)$, which is a free abelian group under addition. If $G = \{(0, \dots, 0)\}$, we say that $X := \{\lambda_1, \dots, \lambda_d\}$ is *multiplicatively independent*. The *rank* of G is the cardinality of a largest multiplicatively independent subset of X . If $\text{rank}(G) = m$, then G has a *basis* $B = \{\mathbf{v}_1, \dots, \mathbf{v}_m\} \subseteq \mathbb{Z}^d$ that is linearly independent over \mathbb{Q} with the property that every $\mathbf{z} \in G$ can be written as an integer linear combination of $\mathbf{v}_1, \dots, \mathbf{v}_m$. If $\lambda_1, \dots, \lambda_d$ are algebraic, we can compute a basis of G using a deep result of Masser [24].

THEOREM 2.4 ([11]). *Given $\lambda_1, \dots, \lambda_d \in \overline{\mathbb{Q}}$, one can compute a basis for $G_M(\lambda_1, \dots, \lambda_d)$.*

Finally, for $\lambda_1, \dots, \lambda_d$ as above, we define the *group of additive relations* as

$$G_A(\lambda_1, \dots, \lambda_d) := \{(k_1, \dots, k_d) \in \mathbb{Z}^d: k_1 \lambda_1 + \cdots + k_d \lambda_d \in \mathbb{Z}\}.$$

Observe that $G_A(\lambda_1, \dots, \lambda_d) = G_M(e^{i2\pi\lambda_1}, \dots, e^{i2\pi\lambda_d})$.

2.4 Linear Recurrence Sequences

A sequence $\langle u_n \rangle_{n=0}^\infty$ is a *linear recurrence sequence* (over \mathbb{Z}) if $u_n \in \mathbb{Z}$ for all $n \in \mathbb{N}$ and there exist $c_1, \dots, c_d \in \mathbb{Z}$ such that $c_d \neq 0$ and

$$u_{n+d} = c_1 u_{n+d-1} + \dots + c_d u_n \quad (1)$$

for all $n \in \mathbb{N}$. We assume that the recurrence (1) is *minimal*, i.e. $c_d \neq 0$. The *characteristic polynomial* of this sequence is $p(x) = x^d - \sum_{i=1}^d c_i x^{d-i}$. Suppose $\langle u_n \rangle_{n=0}^\infty$ is an LRS whose characteristic polynomial has the (distinct) roots $\lambda_1, \dots, \lambda_m \in \overline{\mathbb{Q}}$, called the *characteristic roots* of $\langle u_n \rangle_{n=0}^\infty$. Then there exist unique non-zero polynomials $q_1, \dots, q_m \in \overline{\mathbb{Q}}[x]$ such that

$$u_n = q_1(n)\lambda_1^n + \dots + q_m(n)\lambda_m^n \quad (2)$$

for all $n \in \mathbb{N}$. Eq. (2) is known as the *exponential-polynomial form* of $\langle u_n \rangle_{n=0}^\infty$. For a given characteristic root λ_i , the polynomial q_i is the *coefficient* of λ_i . A characteristic root λ_i is called *simple* if q_i is constant and *dominant* when $|\lambda_i| \geq |\lambda_j|$ for all $1 \leq j \leq m$. We refer the reader to the book [16] for a detailed account of LRS.

Next we give two straightforward lemmas about LRS. First, the exponential-polynomial form (2) immediately implies an exponential upper bound on $|u_n|$, formalised below.

LEMMA 2.5. *Let $\langle u_n \rangle_{n=0}^\infty$ be an LRS, $r, R > 0$ be real algebraic, and suppose $R > |\lambda_i|$ for any characteristic root λ_i of $\langle u_n \rangle_{n=0}^\infty$. We can compute $N \in \mathbb{N}$ such that $|u_n| \leq rR^n$ for all $n \geq N$.*

From (1) it follows that an integer-valued LRS is ultimately periodic modulo any $m \in \mathbb{N}$.

LEMMA 2.6. *Let $\langle u_n \rangle_{n=0}^\infty$ be an LRS and m be a positive integer. We can effectively compute $N, p \in \mathbb{N}$ such that $u_n \equiv u_{n+p} \pmod{m}$ for all $n \geq N$.*

2.5 Schanuel's Conjecture

A set $X = \{\alpha_1, \dots, \alpha_d\}$ of complex numbers is *algebraically independent* over \mathbb{Q} if $p(\alpha_1, \dots, \alpha_d) \neq 0$ for any non-zero polynomial $p \in \mathbb{Q}[x_1, \dots, x_d]$. The *transcendence degree* of X is the size of a largest subset of X that is algebraically independent over \mathbb{Q} . Below we state Schanuel's conjecture, a classical conjecture in transcendental number theory with far-reaching implications [21].

CONJECTURE 2.7 (SCHAUUEL'S CONJECTURE). *If $\alpha_1, \dots, \alpha_d \in \mathbb{C}$ are linearly independent over \mathbb{Q} , then the transcendence degree of $\{\alpha_1, \dots, \alpha_d, \exp(\alpha_1), \dots, \exp(\alpha_d)\}$ is at least d .*

We will use Schanuel's conjecture in two ways. First, consider the structure $\mathbb{R}_{\text{exp}} := \langle \mathbb{R}; <, +, -, \cdot, \exp(\cdot), 0, 1 \rangle$ of real numbers equipped with arithmetic and (real) exponentiation. By the first-order theory of \mathbb{R}_{exp} we mean the set of all well-formed first-order sentences in a suitable language \mathcal{L}_{exp} that are true in \mathbb{R}_{exp} . In [23], Macintyre and Wilkie show that the first-order theory of the structure \mathbb{R}_{exp} is decidable assuming Schanuel's conjecture.

THEOREM 2.8. *Assuming Schanuel's conjecture, given a sentence $\varphi \in \mathcal{L}_{\text{exp}}$, we can decide whether φ holds in \mathbb{R}_{exp} .*

We will also use Schanuel's conjecture to prove the following.

LEMMA 2.9. *Let $\lambda_1, \dots, \lambda_d \in \mathbb{R}_{>1} \cap \overline{\mathbb{Q}}$. Assuming Schanuel's conjecture, a basis for $G_A(1/\log(\lambda_1), \dots, 1/\log(\lambda_d))$ can be computed.*

The proof is in App. A.1.

2.6 Baker's Theorem

By a \mathbb{Q} -affine form we mean $h(x_1, \dots, x_d) = a_0 + \sum_{i=1}^d a_i x_i$, where $a_i \in \mathbb{Q}$ for all $0 \leq i \leq d$. We recall Baker's celebrated theorem on \mathbb{Q} -affine forms in logarithms.

THEOREM 2.10 (THM. 1.6 IN [34]). *Let $\lambda_1, \dots, \lambda_d \in \mathbb{R}_{>0} \cap \overline{\mathbb{Q}}$ be multiplicatively independent. Then the numbers $1, \log(\lambda_1), \dots, \log(\lambda_d)$ are linearly independent over \mathbb{Q} .*

We will use Baker's theorem to prove the following results.

LEMMA 2.11. *Let $d \geq 2$, $\lambda_1, \dots, \lambda_d \in \mathbb{R}_{>1} \cap \overline{\mathbb{Q}}$ be pairwise multiplicatively independent, and suppose*

$$\text{rank}(G_M(\lambda_1, \dots, \lambda_d)) \geq d - 2.$$

Then $1/\log(\lambda_1), \dots, 1/\log(\lambda_d)$ are linearly independent over \mathbb{Q} .

The proof of the lemma above is in App. A.2.

LEMMA 2.12. *Given $\lambda_1, \dots, \lambda_d \in \overline{\mathbb{Q}}$ and $a_0, \dots, a_d \in \mathbb{Q}$, we can effectively determine the sign of $a_0 + \sum_{i=1}^d a_i \log(\lambda_i)$.*

PROOF. By computing the multiplicative relationships among the λ_i using Thm. 2.4, we can rewrite this expression as $b_0 + \sum_{i=1}^e b_i \log(\lambda_i)$. Here, we relabeled the λ_i such that $\lambda_1, \dots, \lambda_e$ is a maximum multiplicatively independent subset of $\lambda_1, \dots, \lambda_d$, and $b_0, \dots, b_e \in \mathbb{Q}$ are explicitly computed. By Thm. 2.10, this expression is 0 if and only if all b_i are 0. If this expression is non-zero, we can compute it up to arbitrary precision and test whether it is positive or not. \square

Multiple effective versions of Baker's theorem exist, which give a lower bound on the magnitude of $\Lambda := h(\log(\rho_1), \dots, \log(\rho_d))$ for a \mathbb{Q} -affine form h , assuming $\Lambda \neq 0$. We use Matveev's version [25] to prove the following in App. A.3.

THEOREM 2.13. *Let $b_1, b_2, c_1, c_2, \rho_1, \rho_2, R_1, R_2$ be positive real algebraic numbers such that $\rho_1 > R_1 > 0$ and $\rho_2 > R_2 > 0$. Then, one can compute $N \in \mathbb{N}$ such that for all $n_1, n_2 \geq N$,*

$$|c_1 \rho_1^{n_1} - c_2 \rho_2^{n_2}| \leq b_1 R_1^{n_1} + b_2 R_2^{n_2} \quad (3)$$

implies that $c_1 \rho_1^{n_1} = c_2 \rho_2^{n_2}$.

2.7 Toric Words

Denote by \mathbb{T} the abelian group \mathbb{R}/\mathbb{Z} , viewed as the interval $[0, 1)$. For $x \in \mathbb{R}$, let $\{x\} := x - \lfloor x \rfloor$ be the fractional part of x . A word $\alpha \in \Sigma^\omega$ is *toric*, written $\alpha \in \mathcal{T}$, if there exist $d > 0$, $\mathbf{s} = (s_1, \dots, s_d) \in \mathbb{T}^d$, a translation $g: \mathbb{T}^d \rightarrow \mathbb{T}^d$ given by

$$(x_1, \dots, x_d) \mapsto (\{x_1 + \delta_1\}, \dots, \{x_d + \delta_d\}) \quad (4)$$

for $\delta_1, \dots, \delta_d \in \mathbb{T}$, and a collection $\mathcal{S} = \{S_b : b \in \Sigma\}$ of subsets of \mathbb{T}^d such that for all $n \in \mathbb{N}$ and $b \in \Sigma$,

$$\alpha(n) = b \Leftrightarrow g^{(n)}(\mathbf{s}) \in S_b. \quad (5)$$

Here $g^{(n)}(\mathbf{s})$ denotes the result of iteratively applying g to \mathbf{s} a total of n times. That is, α is the *coding* (with respect to \mathcal{S}) of the trajectory of the discrete-time dynamical system on \mathbb{T}^d defined by (g, \mathbf{s}) . Observe that $g^{(n)}(\mathbf{s}) = (\{s_1 + n\delta_1\}, \dots, \{s_d + n\delta_d\})$. The word α belongs to the class $\mathcal{T}_{\mathcal{O}}$ of toric words if there exist $d > 0, s_1, \delta_1, \dots, s_d, \delta_d \in \mathbb{T}$ as above and a collection $\{S_b : b \in \Sigma\}$

of *open* subsets of \mathbb{T}^d such that (5) holds for all $n \in \mathbb{N}$ and $b \in \Sigma$. See [7] for a discussion of various subclasses of toric words.

Let $s, \delta \in \mathbb{T}^d$, $G = G_A(\delta)$, and $g: \mathbb{T}^d \rightarrow \mathbb{T}^d$ be as in (4). Define $\mathbb{T}_\delta := \{z \in \mathbb{T}^d : G_A(z) \subseteq G_A(\delta)\} \subseteq \mathbb{T}^d$. The following is a rephrasing of Kronecker's theorem in Diophantine approximation [18]. In the language of dynamical systems, it states that the dynamical system (on \mathbb{T}_δ) obtained by restricting g to \mathbb{T}_δ is *minimal*.

THEOREM 2.14. *The orbit $\langle g^{(n)}(\mathbf{0}) \rangle_{n=0}^\infty$, where $\mathbf{0} = (0, \dots, 0) \in \mathbb{T}^d$, is dense in \mathbb{T}_δ . Moreover, for every open subset O of \mathbb{T}_δ there exist infinitely many $n \in \mathbb{N}$ such that $g^{(n)}(\mathbf{0}) \in O$.*

Let $\mathbf{v}_1, \dots, \mathbf{v}_m \in \mathbb{Z}^d$ be a basis of $G_A(\delta)$, where $m < d$. Write $\mathbf{v}_i = (v_{i,1}, \dots, v_{i,d})$ for $1 \leq i \leq m$ and let $C = \max_i \|\mathbf{v}_i\|_\infty$, noting that $|\mathbf{v}_i \cdot \mathbf{z}| \leq Cd$ for all $\mathbf{z} \in \mathbb{T}^d$. We have that $\mathbf{z} \in \mathbb{T}_\delta$ if and only if $\mathbf{v}_i \cdot \mathbf{z} \in \mathbb{Z}$ for all i , which is equivalent to

$$\bigwedge_{i=1}^m \bigvee_{|k| \leq Cd} \mathbf{v}_i \cdot \mathbf{z} = k.$$

That is, \mathbb{T}_δ , viewed as a subset of \mathbb{R}^d , is an intersection of \mathbb{T}^d with a union of affine subspaces of \mathbb{R}^d with integer parameters. We can now define $\mathbb{T}_{\delta,s} := \{(\{s_1 + z_1\}, \dots, \{s_d + z_d\}) \mid (z_1, \dots, z_d) \in \mathbb{T}_\delta\}$. Applying Thm. 2.14, we obtain the following.

THEOREM 2.15. *The orbit $\langle g^{(n)}(\mathbf{s}) \rangle_{n=0}^\infty$ is dense in $\mathbb{T}_{\delta,s}$, and for every open subset O of $\mathbb{T}_{\delta,s}$ there exist infinitely many $n \in \mathbb{N}$ such that $g^{(n)}(\mathbf{s}) \in O$.*

To prove the first main result of this paper, we will need to show that Acc_α is decidable for certain $\alpha \in \mathcal{T}_O$. The following well-known fact will play an important role in this; See [7] for a proof.

THEOREM 2.16. *Every $\alpha \in \mathcal{T}_O$ is uniformly recurrent.*

2.8 Fourier-Motzkin Elimination

Let $\Phi(x_1, \dots, x_m)$ be a Boolean combination of atomic formulas of the form $h(x_1, \dots, x_m) \sim 0$, where h is a \mathbb{Q} -affine form and \sim is a (strict or non-strict) inequality symbol. Let $1 \leq l \leq m$, and consider the formula $\exists x_1, \dots, x_l \in \mathbb{R} : \Phi(x_1, \dots, x_m)$. Using the Fourier-Motzkin Elimination [14], we can compute a formula $\Psi(x_{l+1}, \dots, x_m) = \bigvee_{j \in J} \bigwedge_{k \in K} h_{j,k}(x_{l+1}, \dots, x_m) \sim_{j,k} 0$ such that

- (a) each $\sim_{j,k}$ is an inequality and $h_{j,k}$ is a \mathbb{Q} -affine form, and
- (b) for all $z_{l+1}, \dots, z_m \in \mathbb{R}$, the sentence

$$\exists x_1, \dots, x_l \in \mathbb{R} : \Phi(x_1, \dots, x_l, z_{l+1}, \dots, z_m)$$

holds if and only if $\Psi(x_{l+1}, \dots, x_m)$ holds.

3 SYNOPSIS OF OUR TECHNIQUES

Our central problem is as follows:

For unary predicates P_1, \dots, P_d , establish the decidability of the MSO theory of $\langle \mathbb{N}; <, P_1, \dots, P_d \rangle$.

In this section, we give a high-level overview of our approach.

The predicates we consider in this paper have arithmetic origins, e.g. $\text{Pow}_2 = \{2^n : n \in \mathbb{N}\}$. In order to exploit their mathematical properties, we view the problem from the equivalent automata-theoretic perspective (Thm. 2.3). Recall the *characteristic word* from Def. 2.2, and let $\alpha = \text{Char}(P_1, \dots, P_d) \in (\{0, 1\}^d)^\omega$. We have that

$\alpha(n) = (b_{n,1}, \dots, b_{n,d})$, where for all $n \in \mathbb{N}$ and $1 \leq i \leq d$, $b_{n,i} = 1$ if $n \in P_i$ and $b_{n,i} = 0$ otherwise. Our problem is thus restated:

For $\alpha = \text{Char}(P_1, \dots, P_d)$, establish the decidability of the automaton acceptance problem Acc_α .

We let $\mathbf{0}$ denote the letter $(0, \dots, 0)$ and express

$$\alpha = \mathbf{0}^{k_0} \alpha(k_0) \mathbf{0}^{k_1} \alpha(k_0 + k_1 + 1) \cdots \mathbf{0}^{k_n} \alpha(k_0 + \dots + k_n + n) \cdots \quad (6)$$

where each letter between the blocks of $\mathbf{0}$'s is not $\mathbf{0}$. Characteristic words defined by a single predicate (e.g. Pow_2 , the powers of 2) are much better studied than those defined by multiple predicates (e.g. $\text{Pow}_2, \text{Pow}_3$). Intuitively, this is because in the latter case, one needs to capture additional element of *interaction* between predicates. To do this, we define *order words*.

Definition 3.1 (Order Word). For unary predicates P_1, \dots, P_d with $\alpha = \text{Char}(P_1, \dots, P_d) \in (\{0, 1\}^d)^\omega$, their order word

$$\alpha' = \text{Ord}(P_1, \dots, P_d) \in (\{0, 1\}^d)^\omega$$

is obtained by deleting all occurrences of $(0, \dots, 0)$ from α .

In compressing the characteristic word α to the order word α' , one only retains partial information, i.e. a particular aspect of the interaction between predicates. Not surprisingly (by Lem. 4.5), $\text{Acc}_{\alpha'}$ always reduces to Acc_α . Remarkably however, under certain circumstances, the interaction captured by order words is the essence of the decision problem Acc_α : For certain well-behaved tuples of predicates, given an automaton \mathcal{A} , one can use the order word α' to recover sufficient information about the interspersed $\mathbf{0}$'s, and hence about α itself, to decide whether \mathcal{A} accepts α .

Let us intuit why this could be so. We rewrite (6) in terms of the order word α' :

$$\alpha = \mathbf{0}^{k_0} \alpha'(0) \cdots \mathbf{0}^{k_n} \alpha'(n) \cdots$$

An automaton \mathcal{A} is crucially finite, and consequently one can compute $K, p > 0$ such that for all $n \geq K$, it cannot distinguish $\mathbf{0}^n$ from $\mathbf{0}^{n+p}$. Provided that k_n is persistently larger than K , it suffices to only keep track of k_n modulo p . We will show that if $\langle \mathbf{0}^{m_n} \rangle_{n=0}^\infty$, where each m_n is indistinguishable from k_n to \mathcal{A} , can be inserted back into α' using a transducer, then Acc_α reduces to $\text{Acc}_{\alpha'}$.

In Sec. 4, we develop the automata-theoretic machinery required to prove that Acc_α (the acceptance problem of the characteristic word) reduces to $\text{Acc}_{\alpha'}$ (the acceptance problem of the order word) assuming the predicates (P_1, \dots, P_d) meet certain technical conditions. In Sec. 5, we invoke Baker's theorem to argue that the predicates defined by LRS with one simple dominant root indeed meet the sufficient conditions identified in Sec. 4.

We devote Sec. 6 to proving decidability of $\text{Acc}_{\alpha'}$ for the order word α' defined by tuples of LRS with one simple dominant root. We do so by showing that α' can be generated as a toric word. Toric words are almost periodic, and it is known that if a word β is *effectively* almost periodic, then Acc_β is decidable (Thm. 4.2). Through number-theoretic arguments, we establish when the order word α' is indeed effectively almost-periodic.

Below is a summary of our strategy to solve our central problem:

- (1) Identify sufficient conditions to reduce Acc_α (the acceptance problem of the characteristic word) to $\text{Acc}_{\alpha'}$ (the acceptance problem of the order word). (Sec. 4)

- (2) Prove that the predicates under consideration meet these conditions. (Sec. 5)
- (3) Generate α' as the trace of a toric dynamical system, and exploit the underlying model to decide $\text{Acc}_{\alpha'}$. (Sec. 6)

In Sec. 7, we follow a similar approach to analyze $\text{Char}(P_1, P_2)$ for $P_1 = \{qn^d : n \in \mathbb{N}\}$ and $P_2 = \{pb^{nd} : n \in \mathbb{N}\}$. The difference is that the underlying dynamical systems are driven by *numeration systems* [22, Chap. 7].

4 PROVING MSO DECIDABILITY

4.1 Classical Results

We recount various classes of words whose automaton acceptance problem is known to be decidable. Recall that a word α is effective if its letters can be effectively computed. Semënov considered the class of (effectively) almost-periodic words in [32].

Definition 4.1. A word $\alpha \in \Sigma^\omega$ is *almost-periodic*, if for every $u \in \Sigma^+$, there exists $R(u) \in \mathbb{N}$ such that the word u either

- (a) does not occur in $\alpha[R(u), \infty)$, or
- (b) occurs in every factor of α of length $R(u)$.

If, moreover, α is effective and a *return time* $R(u)$ as above is computable given u , then α is said to be *effectively almost-periodic*.

THEOREM 4.2 (THEOREM 3 IN [26]). *If $\alpha \in \Sigma^\omega$ is effectively almost-periodic, then Acc_α is decidable.*

Carton and Thomas [12] introduced the class of profinitely ultimately periodic words as a framework to generalise the thematic *contraction methods* of Elgot and Rabin [15].

Definition 4.3. Let Σ be an alphabet.

- (a) A sequence of finite words $\langle u_n \rangle_{n \in \mathbb{N}}$ is *effectively profinitely ultimately periodic* if for any morphism h from Σ^* into a finite monoid M , we can compute N and p such that for all $n \geq N$, $h(u_n) = h(u_{n+p})$.
- (b) An infinite word α is called *effectively profinitely ultimately periodic* if it can be effectively factorised as an infinite concatenation $u_0 u_1 \dots$ of finite non-empty words forming an effectively profinitely ultimately periodic sequence.

THEOREM 4.4. *Let $\alpha \in \Sigma^*$. The problem Acc_α is decidable if and only if α is effectively profinitely ultimately periodic.*

The if direction is due to Carton and Thomas [12], and the converse is due to Rabinovich [28]. Every infinite word is profinitely ultimately periodic, as a close inspection of the use of Ramsey's theorem in Rabinovich's proof reveals. The effectiveness distinguishes words whose automaton acceptance problem is decidable.

A comprehensive class of predicates whose characteristic words ($\alpha \in \{0, 1\}^\omega$) are effectively profinitely ultimately periodic is identified by [12, Thm. 5.2]. This class includes fixed base powers $\text{Pow}_k = \{k^n : n \in \mathbb{N}\}$ as well as k -th powers $\text{N}_k = \{n^k : n \in \mathbb{N}\}$.

4.2 Closure Properties

We now define a few constructs under which the set of infinite words with a decidable automaton acceptance problem is closed.

LEMMA 4.5 (TRANSDUCTION). *Let $\alpha \in \Sigma^\omega$, \mathcal{B} be a deterministic finite transducer with input alphabet Σ and output alphabet Γ , and $\beta = \mathcal{B}(\alpha) \in \Gamma^\omega$. The problem Acc_β reduces to Acc_α .*

PROOF. Given an instance $\mathcal{A} = (Q, q_{\text{init}}, \delta, \mathcal{F})$ of Acc_β , we shall construct \mathcal{A}' whose run on α simulates the run of \mathcal{A} on β . Write $\mathcal{B} = (R, r_{\text{init}}, \sigma)$. The automaton \mathcal{A}' must simulate what \mathcal{B} would do upon reading α , and furthermore, what \mathcal{A} would do upon reading each output block of $\mathcal{B}(\alpha)$. We define the set of states, initial state, and transition function of \mathcal{A}' as

$$\begin{aligned} Q' &= Q \times 2^Q \times R; \\ q'_{\text{init}} &= (q_{\text{init}}, \emptyset, r_{\text{init}}); \\ \delta'((q_1, V_1, r_1), a) &= (q_2, V_2, \sigma_R(r, a)) \end{aligned}$$

such that the invariant $\text{jour}(\sigma_{\Gamma^*}(r, a), q_1) = (q_1, q_2, V_2)$ holds.

Now, express $\beta = u_0 u_1 \dots$ such that \mathcal{B} outputs u_n upon reading the n^{th} letter of α . By construction, the run

$$(q_{\text{init}}, \emptyset, r_{\text{init}})(q_1, V_0, r_1)(q_2, V_1, r_2) \dots$$

of \mathcal{A}' on α corresponds to the run of \mathcal{A} on β , decomposed as the concatenation of journeys $(q_0, q_1, V_0)(q_1, q_2, V_1) \dots$ corresponding to the factorisation $u_0 u_1 \dots$. By Lem. 2.1, a state q is visited infinitely often in \mathcal{A} if and only if a state (q, V, r) with $q \in V$ is visited infinitely often in \mathcal{A}' , and the reduction is thus complete. \square

In fact, we can even use the construction to detect whether β is an infinite word at all: β is infinite if and only if the run of \mathcal{A}' on α visits a state (q, V, r) with $V \neq \emptyset$ infinitely often.

Next, we give our primary closure property. Here we have two words α and β such that α is a ‘‘compressed’’ version of β . We show that Acc_β reduces to Acc_α assuming we can dilate α into (a word equivalent to) β . As expected, Thm. 4.6 further generalises the frameworks of Elgot and Rabin [15] as well as Carton and Thomas [12]. Below, by computing a factorisation $\beta = u_0 u_1 \dots$ we mean giving an algorithm that on every $n \in \mathbb{N}$, returns a finite word u_n such that β is the concatenation of $\langle u_n \rangle_{n=0}^\infty$.

THEOREM 4.6. *Let $\alpha \in \Sigma^\omega$ and $\beta \in \Gamma^\omega$ be such that for any morphism h from Σ^* into a finite monoid M , we can construct a deterministic finite transducer \mathcal{B} (with input alphabet Σ and output alphabet M) and compute a factorisation $u_0 u_1 \dots$ of β such that $\gamma = \mathcal{B}(\alpha) = \gamma(0)\gamma(1)\dots \in M^\omega$ satisfies $h(u_n) = \gamma(n)$ for all $n \in \mathbb{N}$. Then Acc_β reduces to Acc_α .*

PROOF. Given a deterministic finite automaton \mathcal{A} as an instance of Acc_β , we will use the natural morphism h into the finite quotient monoid $M = \Gamma^* / \sim_{\mathcal{A}}$ to reduce it to an instance of Acc_α .

For h, M chosen thus, consider the transducer \mathcal{B} , factorisation $u_0 u_1 \dots$ of β , and word $\gamma = \mathcal{B}(\alpha) \in M^\omega$ from the premise. We have that for all n , $h(u_n) = \gamma(n)$. Since the morphism h maps each word in Γ^* to the equivalence class of words that undertake the same set of journeys, the equivalence class $\gamma(n) \in M$ contains all the information about the journeys that u_n can undertake, i.e. for all $n \in \mathbb{N}$, $q \in Q$, $\text{jour}(u_n, q) = \text{jour}(\gamma(n), q) \in J$.

It is straightforward to construct a transducer \mathcal{B}' such that $\mathcal{B}'(\gamma) = \zeta = j_0 j_1 \dots \in J^\omega$ has the following properties:

- (1) $j_0 = \text{jour}(\gamma(0), q_{\text{init}}) = \text{jour}(u_0, q_{\text{init}}) = (q_{\text{init}}, q_1, V_0)$;
- (2) $j_n = \text{jour}(\gamma(n), q_n) = \text{jour}(u_n, q_n) = (q_n, q_{n+1}, V_n)$ for $n \geq 1$.

We will now show how to decide Acc_β , which is equivalent to deciding whether the run of \mathcal{A} on β visits a state q infinitely often. To that end, observe that the infinite word $\zeta \in J^\omega$ is indeed a

decomposition of the run of \mathcal{A} on β . By Lem. 2.1, the latter visits a state $q \in Q$ infinitely often if and only if $q \in V_n$ of j_n for infinitely many n . This is easily seen to be an instance of Acc_ζ . By Lem. 4.5, Acc_ζ reduces to Acc_γ , which itself reduces to Acc_α . \square

COROLLARY 4.7 (DILATION). *Let $\Sigma = \{1, \dots, b\}$, $\Sigma_0 = \Sigma \cup \{0\}$, and $\alpha \in \Sigma^\omega$. Suppose $\beta \in \Sigma_0^\omega$ and $\beta = 0^{k_0} \alpha(0) 0^{k_1} \alpha(1) \dots$ is such that for any finite monoid M and morphism $h: \Sigma_0^* \rightarrow M$, we can construct a transducer \mathcal{B} with the following property. Its output $\gamma := \mathcal{B}(\alpha) \in M^\omega$ upon reading α satisfies $h(0^{k_n} \alpha(n)) = \gamma(n)$ for all n . Then Acc_β reduces to Acc_α .*

COROLLARY 4.8 (INTERLEAVING). *Let $\alpha \in \Sigma^\omega$, $\langle v_n \rangle_{n \in \mathbb{N}}$ be an effectively profinitely ultimately periodic sequence of words in Σ^* , and $\beta = v_0 \alpha(0) v_1 \alpha(1) \dots \in \Sigma^\omega$. The problem Acc_β reduces to Acc_α .*

PROOF. We will show that for any morphism h into a finite monoid M , we can construct a transducer \mathcal{B} such that $\gamma := \mathcal{B}(\alpha) = \gamma(0)\gamma(1)\dots \in M^\omega$ satisfies $h(v_n \alpha(n)) = \gamma(n)$ for all $n \in \mathbb{N}$. The conclusion then follows immediately from Thm. 4.6.

Since the sequence $\langle v_n \rangle_{n \in \mathbb{N}}$ is effectively profinitely ultimately periodic, we can compute $N, p \in \mathbb{N}$ with $p > 0$ such that $h(v_n) = h(v_{n+p})$ for all $n \geq N$. Define $\gamma(n) = h(v_n \alpha(n)) = h(v_n) \cdot h(\alpha(n))$ for $n < N + p$, and $\gamma(N + qp + r) = h(v_{N+r}) \cdot h(\alpha(N + qp + r))$ for $q > 0$ and $0 < r < p$. It is straightforward to construct a transducer \mathcal{B} that outputs $h(v_0 \alpha(0))h(v_1 \alpha(1))\dots$ upon reading α . \square

4.3 Predicates Corresponding to Functions

Recall that we associate to a tuple of predicates (P_1, \dots, P_d) its characteristic word $\alpha \in \Sigma^\omega$, where $\Sigma = \{0, 1\}^d$ and $\alpha(n)$ records which of the predicates hold for n . The order word α' only records the order in which the predicates hold: it is obtained by deleting all occurrences of $(0, \dots, 0)$ from the characteristic word α .

Definition 4.9. Let \mathcal{F} be the class of strictly increasing functions $f: \mathbb{N} \rightarrow \mathbb{N}$ such that the characteristic word α of the predicate $P = \{f(n) : n \in \mathbb{N}\}$ can be factorised as $0^{k_0} 10^{k_1} 1 \dots$ where $\langle 0^{k_n} \rangle_{n=0}^\infty$ is an effectively profinitely ultimately periodic sequence of words.

The following readily follows from Thm. 4.4.

LEMMA 4.10. *Suppose α is the characteristic word of a predicate $P = \{f(n) : n \in \mathbb{N}\}$ where $f \in \mathcal{F}$. Then Acc_α is decidable.*

THEOREM 4.11 (COMPOSITION). *Let $f_1, \dots, f_d \in \mathcal{F}$. Define functions $g_1, \dots, g_d: \mathbb{N} \rightarrow \mathbb{N}$ such that $g_i = f_1 \circ \dots \circ f_i$. Let $\Sigma = \{0, 1\}^d$, $P_i = \{g_i(n) : n \in \mathbb{N}\}$ for $1 \leq i \leq d$, and $\alpha \in \Sigma^\omega$ be the characteristic word of (P_1, \dots, P_d) . Then Acc_α is decidable.*

PROOF. We will prove the theorem by repeatedly applying Cor. 4.8. Let g_0 be the identity function, $P_0 = \mathbb{N}$, and note that $g_i = g_{i-1} \circ f_i$. Observe that $P_{i+1} = \{g_i(f_{i+1}(n)) : n \in \mathbb{N}\}$ and $P_i = \{g_i(n) : n \in \mathbb{N}\}$. Therefore, $P_1 \supseteq \dots \supseteq P_d$. Let $\alpha^{(0)} = \alpha$, and denote by $\alpha^{(i)}$ the restriction of α to the positions where P_i holds. Let $b_i \in \Sigma$ be the letter whose first i components are 1 and the remaining $d - i$ components are all 0. It is clear that $\alpha^{(i)} \in \{b_i, \dots, b_d\}^\omega$.

Since all functions are strictly increasing, so are their compositions, and we deduce that

$$\alpha^{(i)} = b_i^{k_0} \alpha^{(i+1)}(0) b_i^{k_1} \alpha^{(i+1)}(1) \dots b_i^{k_n} \alpha^{(i+1)}(n) \dots$$

where $\alpha^{(i+1)} \in \{b_{i+1}, \dots, b_d\}^\omega \subset \Sigma^\omega$, $k_0 = f_{i+1}(0)$, and $k_n = f_{i+1}(n) - f_{i+1}(n-1) - 1$. Since $f_{i+1} \in \mathcal{F}$, by Definition 4.9, the sequence $\langle b_i^{k_n} \rangle_{n \in \mathbb{N}}$ is effectively profinitely ultimately periodic. Hence the hypothesis of Cor. 4.8 is satisfied, and $\text{Acc}_{\alpha^{(i)}}$ reduces to $\text{Acc}_{\alpha^{(i+1)}}$. However, $\alpha^{(d)} = b_d^\omega$ and therefore $\text{Acc}_{\alpha^{(d)}}$ is clearly decidable. We conclude that Acc_α is decidable. \square

PROOF OF COR. 1.3. Recall, by [12, Thm. 5.2], $\mathbb{N}_d, \text{Pow}_b \in \mathcal{F}$. On applying the previous theorem with $f_1(n) = n^d$, $f_2(n) = b^n$ we get $g_1(n) = n^d$, $g_2(n) = b^{n^d}$. Thus, Acc_α is decidable for the characteristic word α of the predicates (P_1, P_2) , where $P_1 = \{g_1(n) : n \in \mathbb{N}\}$ and $P_2 = \{g_2(n) : n \in \mathbb{N}\}$. Equivalently, the MSO theory of $\langle \mathbb{N}; <, \mathbb{N}_d, \text{Pow}_{b^d} \rangle$ is decidable. \square

4.4 From Characteristic to Order Words

We will now use our reduction techniques to discuss when the acceptance problem for the characteristic word α is in fact equivalent to that for the order word α' . Note that $\text{Acc}_{\alpha'}$ trivially reduces to Acc_α by Lem. 4.5; we therefore focus on the other direction.

Definition 4.12 (Procylic Predicates). A strictly increasing function $f: \mathbb{N} \rightarrow \mathbb{N}$ is said to be *procylic*³ if the sequence $f(n)$ is effectively ultimately periodic modulo any $m \in \mathbb{N}$, i.e. given any m , there exist computable N, p such that for all $n \geq N$, $f(n+p) \equiv f(n) \pmod{m}$. The corresponding $P = \{f(n) : n \in \mathbb{N}\}$ is called a *procylic predicate*.

Definition 4.13 (Effectively Sparse Predicates). A strictly increasing function $f: \mathbb{N} \rightarrow \mathbb{N}$ is said to be *effectively sparse* if for any $K \in \mathbb{N}$, the inequality $f(n+1) - f(n) \leq K$ has finitely many solutions in n which can moreover be effectively enumerated. The corresponding $P = \{f(n) : n \in \mathbb{N}\}$ is said to be an *effectively sparse predicate*.

Definition 4.14 (Pairwise Effectively Sparse Predicates). Strictly increasing functions $f_1, f_2: \mathbb{N} \rightarrow \mathbb{N}$ are said to be *pairwise effectively sparse* if for any $K \in \mathbb{N}$, the inequality $|f_1(n) - f_2(m)| \leq K$ has finitely many solutions in n, m which can moreover be effectively enumerated. The corresponding $P_1 = \{f_1(n) : n \in \mathbb{N}\}$ and $P_2 = \{f_2(n) : n \in \mathbb{N}\}$ are said to be *pairwise effectively sparse predicates*.

We remark that similar notions of effective sparsity have been considered in literature as early as [31].

THEOREM 4.15 (SPARSE PROCYCLIC PREDICATES). *Let (P_1, \dots, P_d) be a tuple of predicates with characteristic word α and order word α' . Suppose*

- (1) P_1, \dots, P_d are each (a) procylic and (b) effectively sparse;
- (2) each distinct (P_i, P_j) is pairwise effectively sparse.

Then, Acc_α reduces to $\text{Acc}_{\alpha'}$.

PROOF. Let $\mathbf{0} = (0, \dots, 0)$. We observe that

$$\alpha = \mathbf{0}^{k_0} \alpha'(0) \mathbf{0}^{k_1} \alpha'(1) \dots \mathbf{0}^{k_n} \alpha'(n) \dots$$

³This definition is weaker than the notion of (effectively) profinitely ultimately periodic sequences defined by Carton and Thomas [12]. The function f would be effectively profinitely ultimately periodic if, for any morphism h from \mathbb{N} into a finite monoid M , the sequence $\gamma(n) = h(f(n+1) - f(n))$ were effectively ultimately periodic. On the other hand, our definition of procylic functions only considers morphisms into cyclic groups. It is often much easier to show that functions meet the criterion of being procylic rather than being effectively profinitely ultimately periodic.

The above suggests using Cor. 4.7 to prove the theorem. In order to do so, we need to show that for any morphism h into a finite monoid M , we can construct a transducer \mathcal{B} such that $\mathcal{B}(\alpha') = \gamma \in M^\omega$, where $\gamma(n) = h(\mathbf{0}^{k_n} \alpha'(n)) = h(\mathbf{0}^{k_n})h(\alpha'(n))$. The transducer \mathcal{B} outputs $\gamma(n)$ upon reading $\alpha(n)$: it gets the second factor directly from the input. We will show that we can compute the first factor with finite state.

First, note that via a simple pigeonhole argument that exploits the finiteness of monoid M , we can compute $K, m \geq 1$ such that K is a multiple of m , and for all $j \geq K$, $h(\mathbf{0}^j) = h(\mathbf{0}^{j+m})$. In other words, if k_n is sufficiently large, it suffices to only keep track of it modulo m . Now, consider conditions (1b) and (2) of the premise. By the very definition of sparsity, we can compute N such that for all $n \geq N$, $k_n \geq K$. This means that we can compute when k_n is guaranteed to be sufficiently large.

The prefix $\gamma(0) \cdots \gamma(N-1)$ of the output can be hardcoded for the prefix $\alpha'(0) \cdots \alpha'(N-1)$ of the input. It only remains to show how to track k_n modulo m for the infinite suffix. For this, we will use condition (1a) of the premise: that P_1, \dots, P_d are each procyclic. This means that each of f_1, \dots, f_d are effectively ultimately periodic modulo m , i.e. for each i , one only needs finitely many states to evaluate $f_i(n)$, and moreover the state for $f_i(n)$ uniquely determines that for $f_i(n+1)$. As an example, $3^n + 2^n$ modulo 8 follows the pattern 2, 5, 5, 3, 1, 3, 1, 3, \dots , which can be represented by a typical lasso-shaped graph.

The transducer \mathcal{B} thus keeps track of: (a) for each i , what the next occurrence of P_i will be modulo m , and (b) what the occurrence of P_j indicated by the last read letter of $\alpha'(n-1)$ was, modulo m . Upon reading the next letter of $\alpha'(n)$, it can update its record, and appropriately compute k_n , the number of intervening 0's, modulo m . Let this remainder be $r_n < m$. Finally, to write its output, \mathcal{B} simply uses the fact that

$$\gamma(n) = h(\mathbf{0}^{k_n} \alpha'(n)) = h(\mathbf{0}^{k_n})h(\alpha'(n)) = h(\mathbf{0}^{K+r_n})h(\alpha'(n))$$

and we are done. \square

4.5 Normal Words

Recall that a word $\alpha \in \Sigma^\omega$ is *weakly normal* if every $u \in \Sigma^*$ appears infinitely in α .⁴ Normality is usually considered when $\Sigma = \{0, \dots, b-1\}$ is the alphabet of digits and α is the base- b expansion of a real number a . Thus, when $a = \sqrt{2}$ and $b = 10$, $\alpha = 141421356 \cdots$. A number is (weakly) normal in base b if its base- b expansion is a (weakly) normal word. Surveys on normal numbers include Harman [19] and Queffélec [27]. In particular, [19] states the following conjecture.

CONJECTURE 4.16. *An irrational algebraic number α is weakly normal in any integer base $b \geq 2$.*

The strongest result towards this conjecture is due to Adamczewski and Bugeaud [1]. Let $p(n)$ be the number of distinct factors of α of length n . The function p is called the *factor complexity* of α .

⁴In comparison, a *normal word* has the stronger property that each factor u of length n appears with asymptotic frequency $1/|\Sigma|^n$. A real number is *normal* if it is normal in every base b . Constants like $\sqrt{2}$, e , and π are all conjectured to be normal [27].

THEOREM 4.17. *If $b \geq 2$ and α is the base- b expansion of an irrational algebraic number, then*

$$\liminf_{n \rightarrow \infty} \frac{p(n)}{n} = +\infty.$$

We have the following result.

THEOREM 4.18. *If α is weakly normal, then Acc_α is decidable.*

Intuitively, the proof uses the abundance of each factor to guarantee that the set of states visited infinitely often is an entire bottom strongly connected component in the graph induced by the automaton. We defer its technical details to App. A.4.

5 LINEAR RECURRENCE SEQUENCES WITH A SINGLE DOMINANT ROOT

In the upcoming two sections, we prove the first main result of this paper, whose corollaries appeared in the abstract.

THEOREM 5.1. *Consider LRS (over \mathbb{Z}) of the form*

$$u_n^{(i)} = c_i \rho_i^n + \sum_{j=1}^{k_i} p_{i,j}(n) \rho_{i,j}^n,$$

for $1 \leq i \leq d$, such that for all i and j ,

- (1) $c_i, \rho_i, \rho_{i,j} \in \overline{\mathbb{Q}}$ and $p_{i,j} \in \overline{\mathbb{Q}}[x]$,
- (2) $c_i > 0$, $\rho_i > 1$ and $|\rho_{i,j}| < |\rho_i|$, and
- (3) there exist only finitely many pairs $(n, m) \in \mathbb{N}^2$ such that $c_i \rho_i^n = c_j \rho_j^m$.

If we write $P_i \subseteq \mathbb{N}$ for the value set of $\langle u_n^{(i)} \rangle_{n=0}^\infty$, then the MSO theory of $\langle \mathbb{N}; <, P_1, \dots, P_d \rangle$ is decidable assuming Schanuel's conjecture. The decidability is unconditional in either of the following cases:

- (a) If $1/\log(\rho_1), \dots, 1/\log(\rho_d)$ are linearly independent over \mathbb{Q} ;
- (b) If $\text{rank}(G_M(\rho_1, \dots, \rho_d)) \geq d-2$, and ρ_1, \dots, ρ_d are pairwise multiplicatively independent.

This formulation implies Thm. 1.1 from the introduction when ρ_1, \dots, ρ_d are pairwise multiplicatively independent, as conditions (1-3) are immediately met. Suppose ρ_1, \dots, ρ_d are not pairwise multiplicatively dependent. Then $1/\log(\rho_1), \dots, 1/\log(\rho_d)$ are not linearly independent over \mathbb{Q} . We can compute pairwise multiplicatively independent $\lambda_1, \dots, \lambda_m \in \mathbb{N}$ such that for all $1 \leq i \leq d$, $\rho_i = \lambda_{\sigma(i)}^{\mu(i)}$ for some $\sigma(i), \mu(i) \in \mathbb{N}$. Consider $\langle \mathbb{N}; <, \text{Pow}_{\lambda_1}, \dots, \text{Pow}_{\lambda_m} \rangle$. The original characteristic word can be recovered by noting that precisely every $\mu(i)$ -th occurrence of $\text{Pow}_{\lambda_{\sigma(i)}}$ is an occurrence of Pow_{ρ_i} . This can be implemented with a simple transduction; hence by Lem. 4.5, the MSO theory of $\langle \mathbb{N}; <, \text{Pow}_{\rho_1}, \dots, \text{Pow}_{\rho_d} \rangle$ reduces to that of $\langle \mathbb{N}; <, \text{Pow}_{\lambda_1}, \dots, \text{Pow}_{\lambda_m} \rangle$. It remains to observe that if every triple of ρ_1, \dots, ρ_d is multiplicatively dependent, then so is every triple of $\lambda_1, \dots, \lambda_m$.

To prove Thm. 5.1 we will use the framework described in Sec. 3. In the remainder of this section, for $1 \leq i \leq d$, let $\langle u_n^{(i)} \rangle_{n=0}^\infty$ and P_i be as in the statement of Thm. 5.1. Denote by α the characteristic word of (P_1, \dots, P_d) , and by β the order word of (P_1, \dots, P_d) .

Fix $1 \leq i < j \leq d$, and suppose that

$$c_i \rho_i^n = c_j \rho_j^m \tag{7}$$

has solutions $(n_1, m_1), (n_2, m_2) \in \mathbb{N}^2$ in (n, m) . Since $\rho_i, \rho_j > 1$, without loss of generality we can assume $n_1 \leq n_2$ and $m_1 \leq m_2$. Observe that for all $k \in \mathbb{N}$, $(n_1 + k(n_2 - n_1), m_1 + k(m_2 - m_1))$ solves (7). Since (7) has finitely many solutions by assumption, $n_2 = n_1$ and $m_2 = m_1$. Therefore, for every $1 \leq i, j \leq d$, (7) has at most one solution $(n, m) \in \mathbb{N}^2$.

We next define order words obtained from sequences that do not necessarily take exclusively integer values.

Definition 5.2. Let $\langle v_n^{(1)} \rangle_{n=0}^\infty, \dots, \langle v_n^{(d)} \rangle_{n=0}^\infty$ be a family of real-valued, strictly increasing sequences with pairwise disjoint ranges. Further let $Z = \cup_{i=1}^d \{v_n^{(i)} : n \in \mathbb{N}\}$. We define the word

$$\gamma := \text{Ord}(\langle v_n^{(1)} \rangle_{n=0}^\infty, \dots, \langle v_n^{(d)} \rangle_{n=0}^\infty) \in \{1, \dots, d\}^\omega$$

by

$$\gamma(n) = i \Leftrightarrow \exists z \in \langle v_n^{(i)} \rangle_{n=0}^\infty : |\{y \in Z : y < z\}| = n.$$

In this section we prove the following.

THEOREM 5.3. *Let α, β be as above. Acc_α reduces to Acc_β . Furthermore, we can construct positive real algebraic r_1, \dots, r_d such that Acc_β reduces to Acc_ξ , where $\xi = \text{Ord}(\langle r_1 \rho_1^n \rangle_{n=0}^\infty, \dots, \langle r_d \rho_d^n \rangle_{n=0}^\infty)$.*

In Sec. 6 we will show that ξ is effectively almost-periodic and therefore Acc_ξ is decidable. To prove Thm. 5.3, we will need the following lemma, whose main tool is Thm. 2.13.

LEMMA 5.4. *Let $\langle u_n^{(1)} \rangle_{n=0}^\infty, \dots, \langle u_n^{(d)} \rangle_{n=0}^\infty$ be as above. We can compute $N, m_1, \dots, m_d \in \mathbb{N}$ with the following properties.*

- (a) For all $1 \leq i \leq d$, $\langle u_{m_i+n}^{(i)} \rangle_{n=0}^\infty$ is strictly increasing. Moreover, for every $K \in \mathbb{N}$ and $1 \leq i \leq d$, there exists effectively computable L_i such that for all $n \geq L_i$, $u_{n+1}^{(i)} - u_n^{(i)} > K$.
- (b) For all $1 \leq i \leq d$ and $n \in \mathbb{N}$,

$$u_n^{(i)} \geq N \Leftrightarrow n \geq m_i.$$

- (c) $u_{m_1}^{(1)} = N$.
- (d) For all $1 \leq i, j \leq d$, $n_i \geq m_i$ and $n_j \geq m_j$, we have that $c_i \rho_i^{n_i} \neq c_j \rho_j^{n_j}$ and $u_{n_i}^{(i)} \neq u_{n_j}^{(j)}$.
- (e) For $1 \leq i, j \leq d$, $n_i \geq m_i$ and $n_j \geq m_j$,

$$c_i \rho_i^{n_i} > c_j \rho_j^{n_j} \Rightarrow u_{n_i}^{(i)} > u_{n_j}^{(j)}.$$

PROOF. See App. A.5. \square

By Lem. 5.4 (d), each letter of $\alpha[N, \infty)$ is a tuple from $\{0, 1\}^d$ containing at most a single 1. Let $r_i = c_i \rho_i^{m_i}$ for $1 \leq i \leq d$, and $\xi = \text{Ord}(\langle r_1 \rho_1^n \rangle_{n=0}^\infty, \dots, \langle r_d \rho_d^n \rangle_{n=0}^\infty) \in \{1, \dots, d\}^\omega$. By (e), the word $\alpha[N, \infty)$, up to a renaming of variables, is the same as ξ . We can now finalise the proof of Thm. 5.3.

- (1) Each P_i is procyclic. To see this, let $f(n)$ be the n th largest element of P_i and $m \in \mathbb{N}$. We have to show that $\langle f(n) \bmod m \rangle_{n=0}^\infty$ is effectively eventually periodic. By Lem. 5.4 (a), the sequences $\langle u_n^{(i)} \rangle_{n=0}^\infty$ and $\langle f(n) \rangle_{n=0}^\infty$ agree on a suffix that can be effectively determined. It remains to invoke Lem. 2.6.
- (2) By Thm. 6.1 (a), each P_i is effectively sparse.
- (3) Finally, we prove that P_1, \dots, P_d are pairwise effectively sparse. Let $1 \leq i < j \leq d$ and $K \geq 0$. Using Lem. 2.5, compute $R_i, R_j, b_i, b_j > 0$ such that $R_i < \rho_i, R_j < \rho_j$, and for

all $n \geq 0$, $b_i R_i^n > |u_n^{(i)} - c_i \rho_i^n|$ and $b_j R_j^n > |u_n^{(j)} - c_j \rho_j^n + K|$. Then, the triangle inequality and Thm. 2.13 give a $N' \geq 0$ such that for all $n \geq N'$,

$$|u_n^{(i)} - u_n^{(j)}| > |c_i \rho_i^n - c_j \rho_j^n| - b_i R_i^n - b_j R_j^n + K \geq K.$$

By Thm. 4.15, Acc_α reduces to Acc_β . To complete the proof of Thm. 5.3, recall that β can be obtained from ξ through finite modifications, which can be realised by a transducer. Conclude by invoking Lem. 4.5.

6 EFFECTIVE ALMOST PERIODICITY OF THE ORDER WORD

In this section, let $r_i, \rho_i \in \mathbb{R} \cap \overline{\mathbb{Q}}$ with $r_i > 0$ and $\rho_i > 1$ for $1 \leq i \leq d$. Suppose for all $1 \leq i < j \leq d$ and $n, m \in \mathbb{N}$, $r_i \rho_i^n \neq r_j \rho_j^m$. Let $\xi := \text{Ord}(\langle r_1 \rho_1^n \rangle_{n=0}^\infty, \dots, \langle r_d \rho_d^n \rangle_{n=0}^\infty) \in \{1, \dots, d\}^\omega$ as in Def. 5.2. We will prove the following.

THEOREM 6.1.

- (a) The word ξ is almost-periodic.
- (b) Assuming Schanuel's conjecture, ξ is effectively almost-periodic.
- (c) If $1/\log(\rho_1), \dots, 1/\log(\rho_d)$ are linearly independent over $\overline{\mathbb{Q}}$, then ξ is unconditionally effectively almost-periodic.
- (d) If $\text{rank}(G_M(\rho_1, \dots, \rho_d)) \geq d - 2$, and ρ_1, \dots, ρ_d are pairwise multiplicatively independent, then ξ is unconditionally effectively almost-periodic.

This result, together with Thm. 5.3, will prove Thm. 5.1. By scaling and reordering the d sequences, **we can without loss of generality assume that $1 < \rho_1 \leq \rho_2 \leq \dots \leq \rho_d$ and $r_1 = 1$** . Let

$$V = \{\log(r_i \rho_i^n) : 1 \leq i \leq d, n \in \mathbb{N}\}$$

and $\langle v_n \rangle_{n=0}^\infty$ be the ordering of V with $v_0 < v_1 < \dots$. Observe that

$$\xi = \text{Ord}(\langle \log(r_1 \rho_1^n) \rangle_{n=0}^\infty, \dots, \langle \log(r_d \rho_d^n) \rangle_{n=0}^\infty).$$

Example 6.2. Consider $r_1 = 1, \rho_1 = 2, r_2 = 9, \rho_2 = 3$. Then



$$2^0 < 2^1 < 2^2 < 2^3 < 9 \cdot 3^0 < 2^4 < 9 \cdot 3^1 < 2^5 < \dots$$

and hence $\xi = 11112121121 \dots$. We have $v_0 = 0, v_1 = \log(2), v_2 = 2 \log(2), v_3 = 3 \log(2), v_4 = \log(9)$, and so on.

Our strategy to prove Thm. 6.1 will be to show that ξ has a suffix that belongs to the class \mathcal{T}_0 of toric words; recall from Sec. 2.7 that words in \mathcal{T}_0 are uniformly recurrent and hence almost-periodic. To prove effective almost-periodicity, we will deploy number theory either through Baker's theorem, or Schanuel's conjecture.

Continuing Ex. 6.2, let $a_n = n \log(2)$ and $b_n = \log(9) + n \log(3)$. Figure 1 illustrates a way to generate ξ . We start at the point $(0, 0)$ and follow the line $y = x$. Every time a vertical line $x = a_n$ for some n is hit, we write a **1**. When we hit a horizontal line $y = b_n$ for some n , we write **2**. If we discard the first three characters of ξ , we obtain a *cutting sequence* (equivalently, a *billiard word*), illustrated in Fig. 2. (See [17, Chap. 4.1.2] and [3, 4] for more on billiard words.) Figure 2 is obtained from Fig. 1 by a translation and a scaling. In Fig. 2, we start at the point $(0, y)$, where $0 < y < 1$, and follow the dashed line which has slope $\log(2)/\log(3)$. When we hit a line $x = n$ for $n \in \mathbb{N}$, we write **1**; When we hit $y = n$, we write **2**.

We will not directly use the fact that a suffix of ξ is a cutting sequence, but combinatorial properties of such sequences can be


 Figure 1: Generating ξ .

 Figure 2: The suffix $\xi[3, \infty)$ as a cutting sequence.

used to prove a weaker version of Thm. 6.1 (c); see Sec. 6.1. We note that cutting sequences generated by a line on the plane with irrational slope (as in Fig. 2) are exactly the Sturmian words [22].

We continue our proof of Thm. 6.1. Define $z_n = \frac{v_n}{\log(\rho_1)}$, and observe that $\langle z_n \rangle_{n=0}^\infty$ is strictly increasing. Let $\sigma: \{z_n : n \in \mathbb{N}\} \rightarrow \{1, \dots, d\}$ be such that $\xi(n) = \sigma(z_n)$. Observe that for every n , there exists $m \in \mathbb{N}$ such that $z_m = n$ and $\sigma(z_m) = 1$. For this value of m , $v_m = \log(r_1 \rho_1^m) = n \log(\rho_1)$. In Ex. 6.2, $z_0 = 0$, $z_1 = 1$, $z_2 = 2$, $z_3 = 3$, $z_4 = \log(9)/\log(2) \approx 3.17$, $z_5 = 4$, $z_6 = \log(9 \cdot 3)/\log(2) \approx 4.75$, $z_7 = 5$, $z_8 = 6$, and so on.

We will next factorise $\xi = w_- w_0 w_1 \dots$. Intuitively, for $n \in \mathbb{N}$, the finite word w_n contains the labels of all terms of $\langle z_n \rangle_{n=0}^\infty$ (obtained by applying σ) that lie in the interval $[n, n+1)$. Formally, let $\langle k_n \rangle_{n=0}^\infty$ be the sequence over \mathbb{N} such that for all $n, m \in \mathbb{N}$, $n \leq v_m < n+1$ if and only if $k_n \leq m < k_{n+1}$. Then $w_- = \xi[0, k_0)$ and $w_n = \xi[k_n, k_{n+1})$ for all $n \in \mathbb{N}$. Consider $n \in \mathbb{N}$. As argued earlier, there exists $m \in \mathbb{N}$ such that $z_m = n$ and $\sigma(z_m) = 1$. That is, the first letter of w_n for every n is 1. Moreover, in each such w_n the letter 1 occurs exactly once. In Ex. 6.2, w_- is empty, $w_0 = w_1 = w_2 = 1$, $w_3 = 12$, $w_4 = 12$, $w_5 = 1$, and so on.

Let N be the smallest integer n such that every letter of $\{1, \dots, d\}$ appears in $w_0 \dots w_N \in \{1, \dots, d\}^*$. In Ex. 6.2, $N = 3$. Further let Σ be the set of all finite patterns over $\{1, \dots, d\}$ that start with the letter 1 and contain at most one instance of i for all $i \in \{1, \dots, d\}$. Define $\beta \in \Sigma^\omega$ by $\beta(n) = w_{N+n}$. In our example, $\beta = (12)(12)(1) \dots$. Observe that the suffix $w_N w_{N+1} \dots$ of ξ is the image of β under the application of the morphism $\mu: \Sigma^* \rightarrow \{1, \dots, d\}^*$ defined by $\mu(w) = w(0) \dots w(|w| - 1)$ for $w \in \Sigma^*$. Since effectively almost-periodic words are closed under applications of morphisms (provided that the image word is also infinite) and finite modifications [26], the word β is effectively almost-periodic if and only if ξ is. We will next show that β is toric with $\beta \in \mathcal{T}_O$. Recall that $\{x\} := x - \lfloor x \rfloor$ denotes the fractional part of x .

THEOREM 6.3. For $2 \leq i \leq d$, let $\delta_i = \frac{\log(\rho_1)}{\log(\rho_i)} \in \mathbb{T}$ and

$$s_i = \left\{ \frac{(N+1) \log(\rho_1) - \log(r_i)}{\log(\rho_i)} \right\} \in \mathbb{T}.$$

Then β is the toric word generated by $\delta = (\delta_2, \dots, \delta_d) \in \mathbb{T}^{d-1}$ and $s = (s_2, \dots, s_d) \in \mathbb{T}^{d-1}$, as well as a collection of open subsets of \mathbb{T}^{d-1} defined by linear inequalities (in variables x_2, \dots, x_d) of the form $x_i/\delta_i < x_j/\delta_j$ or $x_i/\delta_i \sim 1$, where $\sim \in \{>, <\}$ and $2 \leq i, j \leq d$.

PROOF. For $2 \leq i \leq d$ and $n \in \mathbb{N}$, let

$$L(n, i) = \left\{ \frac{\log(r_i \rho_i^k)}{\log(\rho_1)} : k \in \mathbb{N} \right\} \cap (-\infty, N+n+1]$$

and $\Delta(n, i)$ be the smallest distance from $N+n+1$ to an element of $L(n, i)$. Intuitively, $L(n, i)$ is the “downward distance” from $N+n+1$ to the lattice $\{\log(r_i \rho_i^k)/\log(\rho_1) : k \in \mathbb{N}\}$. By construction of N , the value of $\Delta(n, i)$ is well-defined and finite for every n and i .

Let $b = b_0 b_1 \dots b_m \in \Sigma$, where $b_i \in \{1, \dots, d\}$ for all i and $b_0 = 1$. By definition, $\beta(n) = b$ if and only if $w_{N+n} = b$, which is the case if and only if the following hold.

(*) There exist $k_1, \dots, k_m \in \mathbb{N}$ such that

$$N+n \leq \frac{\log(r_{b_1} \rho_{b_1}^{k_1})}{\log(\rho_1)} < \dots < \frac{\log(r_{b_m} \rho_{b_m}^{k_m})}{\log(\rho_1)} < N+n+1.$$

(**) For every $k \in \mathbb{N}$ and $i \in \{2, \dots, d\}$ not appearing in b ,

$$\frac{\log(r_i \rho_i^k)}{\log(\rho_1)} \notin [N+n, N+n+1).$$

As discussed earlier, for every $t \in \mathbb{N}$ there exist $l \in \mathbb{N}$ such that

$$z_l = t = \log(r_1 \rho_1^t)/\log(\rho_1).$$

Since no two distinct terms of $\langle z_n \rangle_{n=0}^\infty$ are equal, for every $i \neq 1$ and $k \in \mathbb{N}$, $\log(r_i \rho_i^k)/\log(\rho_1) \notin \mathbb{N}$. Thus in (*) and (**) we can replace non-strict inequalities with strict ones, and vice versa.

Next, observe that (*) is equivalent to

$$\Delta(n, b_m) < \dots < \Delta(n, b_1) < 1. \quad (8)$$

Similarly, (**) holds if and only if for every $i \in \{2, \dots, d\}$ not appearing in b ,

$$\Delta(n, i) > 1. \quad (9)$$

Since for all i and $k \in \mathbb{N}$,

$$\frac{\log(r_i \rho_i^k)}{\log(\rho_1)} = \frac{\log(r_i)}{\log(\rho_1)} + k \frac{\log(\rho_i)}{\log(\rho_1)},$$

we have that

$$\Delta(n, i) = \frac{\log(\rho_i)}{\log(\rho_1)} \cdot \left\{ \frac{N+n+1 - \log(r_i)/\log(\rho_1)}{\log(\rho_i)/\log(\rho_1)} \right\} = \frac{1}{\delta_i} \{s_i + n\delta_i\}.$$

Consider the dynamical system on \mathbb{T}^{d-1} given by (g, s) , where $g(y_2, \dots, y_d) = (\{y_2 + \delta_2\}, \dots, \{y_d + \delta_d\})$. Recall from Sec. 2.7 that for $n \in \mathbb{N}$, $g^{(n)}(s) = (\{s_2 + n\delta_2\}, \dots, \{s_d + n\delta_d\})$. Considering (8) and (9), we are led to defining, for $b = b_0 b_1 \dots b_m \in \Sigma$, $S_b \subseteq \mathbb{T}^{d-1}$ as the (open) set of all $(x_2, \dots, x_d) \in \mathbb{T}^{d-1}$ such that

$$\frac{x_{b_m}}{\delta_{b_m}} < \dots < \frac{x_{b_1}}{\delta_{b_1}} < 1 \quad \wedge \quad \bigwedge_{\substack{1 \leq j \leq d \\ j \neq b_1, \dots, b_m}} \frac{x_j}{\delta_j} > 1. \quad (10)$$

12	1
132	13
	123

Figure 3: The torus for $\rho_1 = 2$, $\rho_2 = 3$, and $\rho_3 = 5$.

With this definition, for all $b \in \Sigma$ and $n \in \mathbb{N}$, $\beta(n) = b$ if and only if $g^{(n)}(s) \in S_b$. \square

In the remainder of this section, let $s = (s_2, \dots, s_d) \in \mathbb{T}^{d-1}$ and $\delta := (\delta_2, \dots, \delta_d) \in \mathbb{T}^{d-1}$, and $g: \mathbb{T}^{d-1} \rightarrow \mathbb{T}^{d-1}$ be defined as above. Figure 3 illustrates the target sets $\{S_b : b \in \Sigma\}$ constructed in Thm. 6.3 for the sequences $\langle 2^n \rangle_{n=1}^\infty$, $\langle 3^n \rangle_{n=1}^\infty$, and $\langle 5^n \rangle_{n=1}^\infty$. Figure 3 can also be viewed as follows. Consider a dynamical system on \mathbb{N} that starts at 2 and at each step jumps to the next power of 2. At each step, a letter from $\{1, 12, 13, 123, 132\}$ is written depending on whether the point jumped over a power of 3 or a power of 5 in the last step (and in what order). The exact letter to be written is determined by keeping track of the fractional part of $\log_3(2^n)$ and $\log_5(2^n)$; this gives rise to the linear inequalities defining the open sets depicted in Fig. 3.

Since $\beta \in \mathcal{T}_O$, it is uniformly recurrent (Sec. 2.7). As mentioned earlier, the suffix $w_N w_{N+1} \dots$ of ξ is the image of β under a morphism, and hence is almost-periodic by [26, Sec. 3]. It follows that ξ is almost-periodic; **this proves Thm. 6.1 (a)**. We next analyse effective almost periodicity of β , which implies effective almost periodicity of ξ . We will need the following lemma, whose proof can be found in App. A.6.

LEMMA 6.4. *Let α be a uniformly recurrent word for which $\alpha(n)$ can be effectively determined given n . Then the word α is effectively almost-periodic if and only if we can decide occurrence of a given finite word w in α .*

We next study how to decide whether given $w \in \Sigma^*$ occurs in β .

LEMMA 6.5. *Let $w \in \Sigma^*$. There exists an open subset $S_w \subseteq \mathbb{T}^{d-1}$ with the following property. For all $n \in \mathbb{N}$, the pattern w occurs in β at the position n if and only if $g^{(n)}(s) \in S_w$. Furthermore, we can compute a representation of S_w as a Boolean combination of inequalities of the form*

$$h(x_2/\delta_2, \dots, x_d/\delta_d, 1/\delta_2, \dots, 1/\delta_d) \sim 0, \quad (11)$$

where h is a \mathbb{Q} -affine form and \sim is an inequality symbol.

PROOF. See App. A.7. \square

We next prove Thm. 6.1 (c). Suppose $1/\log(\rho_1), \dots, 1/\log(\rho_d)$ are linearly independent over \mathbb{Q} . Then for all $c_2, \dots, c_d, k \in \mathbb{Q}$,

$$\sum_{i=2}^d c_i \delta_i = k \Leftrightarrow \frac{c_2}{\log(\rho_2)} + \dots + \frac{c_d}{\log(\rho_d)} = \frac{k}{\log(\rho_1)}$$

$$\Leftrightarrow k, c_2, \dots, c_d = 0.$$

Hence $G_A(\delta)$ is the trivial group, $\mathbb{T}_\delta = \mathbb{T}^{d-1}$, and by Kronecker's theorem (Thm. 2.14), $\langle g^{(n)}(s) \rangle_{n=0}^\infty$ is dense in \mathbb{T}^{d-1} . Therefore, a pattern $w \in \Sigma^*$ occurs in β if and only if $S_w \neq \emptyset$. As shown in

Lem. 6.5, to decide whether $S_w \neq \emptyset$ we have to decide the truth of $\Psi := \exists x_2, \dots, x_d: \Phi(x_2, \dots, x_d)$ where Φ is a formula of the form

$$\bigvee_{j \in J} \bigwedge_{k \in K} h_{j,k}(x_2/\delta_2, \dots, x_d/\delta_d, 1/\delta_2, \dots, 1/\delta_d) \sim_{j,k} 0.$$

Recall that each $h_{j,k}$ is a \mathbb{Q} -affine form. Hence Ψ is equivalent to $\exists \tilde{x}_2, \dots, \tilde{x}_d: \Phi(\tilde{x}_2, \dots, \tilde{x}_d)$. Applying Fourier-Motzkin Elimination, we can compute finitely many \mathbb{Q} -affine forms $h_{l,m}$ and an inequality symbols $\sim_{l,m}$ such that Ψ is true if and only if

$$\bigvee_{l \in L} \bigwedge_{m \in M} h_{l,m}(1/\delta_2, \dots, 1/\delta_d) \sim_{l,m} 0.$$

Recall that $\frac{1}{\delta_i} = \frac{\log(\rho_i)}{\log(\rho_1)}$. For $h(x_2, \dots, x_d) := c_1 + c_2 x_2 + \dots + c_d x_d$,

$$h(1/\delta_2, \dots, 1/\delta_d) = \frac{1}{\log(\rho_1)} (c_1 \log(\rho_1) + \dots + c_d \log(\rho_d)).$$

Hence for all l, m , whether $h_{l,m}(1/\delta_2, \dots, 1/\delta_d) \sim_{l,m} 0$ can be decided using Baker's theorem (Lem. 2.12). Thus, under the assumption that $1/\log(\rho_1), \dots, 1/\log(\rho_d)$ are linearly independent over \mathbb{Q} , we can decide whether a given pattern $w \in \Sigma^*$ occurs in β . We conclude that β and hence ξ are effectively almost periodic. **This proves Thm. 6.1 (c). To prove Thm. 6.1 (d), recall Lem. 2.11 and invoke Thm. 6.1 (c).**

It remains to prove Thm. 6.1 (b). Assuming Schanuel's conjecture, we can compute a basis of $G_A(1/\log(\rho_1), \dots, 1/\log(\rho_d))$ (Lem. 2.9). Hence we can compute an \mathbb{R}_{exp} formula defining the compact $\mathbb{T}_{\delta,s} \subseteq \mathbb{T}^{d-1}$ in which $\langle f^{(n)}(s) \rangle_{n=0}^\infty$ is dense (Sec. 2.7). Recall from Sec. 2.7 that a pattern w occurs in β if and only if $S_w \cap \mathbb{T}_{\delta,s} \neq \emptyset$, which can be effectively verified using a decision procedure for the first-order theory of \mathbb{R}_{exp} . Hence β and ξ are effectively almost-periodic assuming Schanuel's conjecture. **This proves Thm. 6.1 (b).**

6.1 Applying the Theory of Cutting Sequences

Let $\xi = \text{Ord}(\langle r_1 \rho_1^n \rangle_{n=0}^\infty, \dots, \langle r_d \rho_d^n \rangle_{n=0}^\infty)$ be as above. As hinted earlier, it can be shown that ξ has a suffix that is the cutting sequence generated by the line $\{(s_1 + t/\log(\rho_1), \dots, s_d + t/\log(\rho_d)) : t \geq 0\}$, where $s_i \in [0, 1)$ for all i . As in Sec. 4.5, we write $p(n)$ for the number of distinct factors of ξ of length n ; the function p is the factor complexity of ξ . The factor complexity of cutting sequences has been extensively studied, and in many cases, an exact formula for $p(n)$ is known. We give an overview of results in this direction.

- (i) If $d = 2$ and $\log(\rho_1)/\log(\rho_2)$ is irrational, then ξ is a Sturmian word and therefore $p(n) = n + 1$. See, e.g. [2, Chap. 10.5].
- (ii) By [3], if $d = 3$, and $1/\log(\rho_1), 1/\log(\rho_2), 1/\log(\rho_3)$ as well as $\log(\rho_1), \log(\rho_2), \log(\rho_3)$ are linearly independent over \mathbb{Q} , then $p(n) = n^2 + n + 1$.
- (iii) For arbitrary $d > 0$, Bedaride [6] gives an exact formula for $p(n)$ assuming $1/\log(\rho_1), \dots, 1/\log(\rho_d)$ as well as every triple $\log(\rho_i), \log(\rho_j), \log(\rho_k)$ for pairwise distinct i, j, k are linearly independent over \mathbb{Q} . This generalises the well-known result [4] of Baryshnikov.

Going back to our word ξ , let w be a finite pattern of length n , and suppose we know the value of $p(n)$. Then we can decide whether w occurs (as required by Lem. 6.4) in ξ by just reading prefixes of ξ until we have seen $p(n)$ distinct factors of length n . Using

this approach, we can prove that the word ξ is effectively almost-periodic under the assumption of (iii). Note, however, that this result is strictly weaker than Thm. 6.1 (b). Consider, for example, $\rho_1 = 2$, $\rho_2 = 3$ and $\rho_3 = 6$. By Lem. 2.11, $1/\log(\rho_1), \dots, 1/\log(\rho_3)$ are linearly independent over \mathbb{Q} , but $\log(\rho_1), \dots, \log(\rho_3)$ are not.

6.2 Proof of Theorem 5.1

We can now combine everything we have shown so far to prove Thm. 5.1. For $1 \leq i \leq d$, let $\langle u_n^{(i)} \rangle_{n=0}^\infty$ for $1 \leq i \leq d$ be as in the statement of Thm. 5.1 with the value set P_i . Further let α be the characteristic word of $\langle P_1, \dots, P_d \rangle$, and recall that the MSO theory of $\langle \mathbb{N}; <, P_1, \dots, P_d \rangle$ is decidable if and only if Acc_α is decidable. Applying Thm. 5.3, we can construct r_1, \dots, r_d such that Acc_α reduces to Acc_ξ , where $\xi = \text{Ord}(\langle r_1 \rho_1^n \rangle_{n=0}^\infty, \dots, \langle r_d \rho_d^n \rangle_{n=0}^\infty)$. Applying Thm. 6.1, we obtain conditions under which ξ is effectively almost-periodic. It remains to recall from Thm. 4.2 that Acc_ξ is decidable if ξ is effectively almost-periodic.

7 MSO OF LRS AND NORMAL NUMBERS

In this section, we discuss a second class of LRS that give rise to interesting MSO theories; we will show that these are intimately connected to base- b expansions of certain algebraic numbers. In particular, we will show that the base- b expansion of $\sqrt[d]{p/q}$ is intrinsic to the pair of predicates $\{qn^d : n \in \mathbb{N}\}$ and $\{pb^{nd} : n \in \mathbb{N}\}$. That is, the binary expansion of $\sqrt[3]{1/27} = 1/3$ underlies the pair of predicates $\{27n^3 : n \in \mathbb{N}\}$ and $\{8^n : n \in \mathbb{N}\}$, while the binary expansion of $\sqrt[5]{5}$ underlies the pair $\{n^3 : n \in \mathbb{N}\}$ and $\{5 \cdot 8^n : n \in \mathbb{N}\}$. The dynamical systems at play in this section differ from the ones we considered previously: they are defined by *numeration systems* [22, Chap. 7] as opposed to translations on a torus (Sec. 6 and 2.7).

We begin by considering the case where $\sqrt[d]{p/q}$ is rational, which implies that its base- b expansion is ultimately periodic for any $b \geq 2$. The following is a generalisation of Cor. 1.3; see App. A.8 for the proof, which is a simple application of Thm. 4.11.

THEOREM 7.1. *Let $b, d \geq 2$ and $p, q \geq 1$ be integers such that $\sqrt[d]{p/q}$ is rational. Let $\alpha \in \{0, 1\}^\omega$ be the characteristic word of $\langle P_1, P_2 \rangle$, where $P_1 = \{qn^d : n \in \mathbb{N}\}$ and $P_2 = \{pb^{nd} : n \in \mathbb{N}\}$. Then the problem Acc_α is decidable.*

The case where $\sqrt[d]{p/q}$ is irrational is more involved.

THEOREM 7.2. *Let b, d, p, q be positive integers such that $\sqrt[d]{p/q}$ is irrational. Furthermore, let*

- (1) $\alpha \in \{0, 1\}^\omega$ be the characteristic word of $\langle P_1, P_2 \rangle$, where $P_1 = \{qn^d : n \in \mathbb{N}\}$ and $P_2 = \{pb^{nd} : n \in \mathbb{N}\}$;
- (2) $\beta \in \{0, 1, \dots, b-1\}^\omega$ be the infinite string of digits in the base- b expansion of $\eta = \sqrt[d]{p/q}$ and
- (3) γ be the order word corresponding to α , i.e. the word obtained by deleting all occurrences of $(0, 0)$ from α .

Then the problems Acc_α , Acc_β , and Acc_γ are Turing-equivalent.

PROOF. We will prove the theorem by showing:

- (1) Acc_β reduces to Acc_α .
- (2) Acc_α reduces to Acc_γ .
- (3) Acc_γ reduces to Acc_β .

Part (1): Acc_β reduces to Acc_α . By construction (except for an easily computable finite prefix), we have the invariant

$$\beta(n) = \lfloor \eta b^n \rfloor \bmod b, \quad \beta(n) \in \{0, \dots, b-1\}.$$

For example, if $p = 2$, $q = 1$, $b = 10$, and $d = 2$, then $\eta = \sqrt{2} = 1.4142 \dots$ and $\beta(0) = 1, \beta(1) = 4, \beta(2) = 1$ etc. This observation accounts for one reduction of the Turing equivalence.

We will prove the claim by constructing a deterministic finite transducer \mathcal{B} such that $\beta = \mathcal{B}(\alpha)$ and applying Lem. 4.5. The states of \mathcal{B} are $R = \{0, \dots, b-1\}$ with initial state $b-1$. The transducer moves from state q to state $q+1 \bmod b$ if it reads $(1, 0)$ and does not move otherwise. The transducer outputs its current state q on reading $(0, 1)$ and otherwise it outputs the empty word. Thus, the letter $(0, 0)$ has no effect on the state or output of the transducer. By construction, the transducer keeps count of the number of occurrences of P_1 modulo b , and outputs this count on encountering an occurrence of P_2 . Hence its n th output will be $b_n \bmod b$, where $b_n := |\{m \in \mathbb{N} : qm^d < pb^{nd}\}| - 1$. It remains to observe that

$$b_n = |\{m \in \mathbb{N} : m < b^n \cdot \sqrt[d]{p/q}\} - 1| \equiv \lfloor \eta b^n \rfloor \pmod{b}.$$

Part (2): Acc_α reduces to Acc_γ . This reduction requires a more refined understanding of the word α . We readily observe that $(1, 1)$ does not occur in α . If $qm^d = pb^{nd}$, then $m = \eta b^n$. As η is irrational, m and b^n cannot both be integers. Thus, α only contains $(0, 0)$, $(0, 1)$ and $(1, 0)$.

Consider the order word $\gamma \in \{(1, 0), (0, 1)\}^\omega$ of $\langle P_1, P_2 \rangle$.

$$\alpha = (0, 0)^{n_0} \gamma (0, 0)^{n_1} \gamma (1) \dots$$

We apply Thm. 4.15 to show that in the case of $\langle P_1, P_2 \rangle$, the acceptance problem of the characteristic word reduces to that of the order word. Condition (1) of the premise, i.e. that P_1 and P_2 are pro-cyclic and effectively sparse, are easily seen to hold. We have also established that P_1 and P_2 can never hold simultaneously. To argue Condition (2), i.e. that P_1, P_2 are pairwise effectively sparse, we apply the following result of Schinzel and Tijdeman [30].

LEMMA 7.3 (SCHINZEL AND TIJDEMAN). *For every $N \geq 1$, the equation $|qn^d - pb^{md}| = N$ has finitely many solutions (n, m) that can be effectively enumerated.*

Hence the claim follows from Thm. 4.15: Acc_α of the characteristic word reduces to Acc_γ of the order word.

Part (3): Acc_γ reduces to Acc_β . Note that γ itself can be written as

$$\gamma = (1, 0)^{m_0} (0, 1) (1, 0)^{m_1} (0, 1) \dots$$

We now reduce Acc_γ to Acc_β using Cor. 4.7. We remark that despite the differing underlying arithmetic, the key automata-theoretic ideas are similar to the proof of Thm. 4.15. We record the invariants:

$$S_k = \sum_{i=0}^k m_i = \lfloor \eta b^k \rfloor$$

$$\beta_k \equiv \lfloor \eta b^k \rfloor \pmod{b}$$

$$bS_k \leq S_{k+1} < b(S_k + 1)$$

$$S_{k+1} = bS_k + \beta_{k+1} = S_k + m_{k+1}.$$

Define

$$\gamma' = \perp^{m_0} \beta(0) \perp^{m_1} \beta(1) \cdots$$

By Lem. 4.5, it is easy to see that Acc_γ reduces to $\text{Acc}_{\gamma'}$ via a straightforward transduction, which is in fact a homomorphism. It thus suffices to reduce $\text{Acc}_{\gamma'}$ to Acc_β .

We denote $S_k = \lfloor \eta b^k \rfloor$ and observe that $bS_k \leq S_{k+1} < b(S_k + 1)$ (one can check the identity $b\lfloor x \rfloor \leq \lfloor bx \rfloor < b(\lfloor x \rfloor + 1)$). Thus, $S_{k+1} = bS_k + \beta_{k+1} = S_k + m_{k+1}$. Given β and m_0 , we can use the above properties to track S_k , and hence m_k modulo t for any integer t . This is useful, because: (1) it is easy to observe that $\langle m_k \rangle_{k \in \mathbb{N}}$ is a strictly increasing sequence, and that we can effectively compute j such that $m_k \geq N$ for all $k \geq j$; (2) for any morphism h into a finite monoid M , there exist effective N, t such that for all $n \geq N$, $h(\perp^n) = h(\perp^{n+t})$.

Using these observations, given any morphism h into a finite monoid M , we can construct a transducer \mathcal{B} such that $\mathcal{B}(\beta) = \mu \in M^\omega$. We recall that we take the j^{th} factor of γ' to be $\perp^{m_j} \beta(j)$. On reading $\beta(j)$, \mathcal{B} outputs μ_j , with the property that $h(\perp^{m_j} \beta(j)) = \mu_j$ for all $j \in \mathbb{N}$. The hypothesis of Cor. 4.7 is met, and $\text{Acc}_{\gamma'}$ thus reduces to Acc_β . \square

By Conj. 4.16, it is expected that $\sqrt[d]{p/q}$ is a weakly normal number in base b when it is irrational. Hence, if the conjecture holds, Acc_β is decidable by Thm. 4.18. Applying Theorem 7.2, both Acc_α and Acc_γ are decidable assuming Conj. 4.16.

We can slightly simplify the special case of Thm. 7.2 where $p = b, q = 1, d = 2$ to the procyclic functions n^2 and b^n ; see App. A.9 for the proof, which closely mirrors that of Thm. 7.2. On applying this result with $b = 2$, we get that the MSO theory of $\langle \mathbb{N}; <, N_2, \text{Pow}_2 \rangle$ is Turing-equivalent with Acc_β , where β is the binary expansion of $\sqrt{2}$, as stated in the abstract.

THEOREM 7.4. *Let $b \geq 2, P_1 = N_2, P_2 = \text{Pow}_b$, and $\Sigma = \{0, 1\}^2$. Further let $\alpha \in \Sigma^\omega$ be the characteristic word of (P_1, P_2) and $\beta \in \{0, \dots, b-1\}^\omega$ be the base- b expansion of \sqrt{b} . Then the problems Acc_α and Acc_β are Turing-equivalent.*

ACKNOWLEDGMENTS

Toghrol Karimov, Joël Ouaknine, and Mihir Vahanwala were supported by DFG grant 389792660 as part of TRR 248 (see <https://perspicuous-computing.science>). Joël Ouaknine is also affiliated with Keble College, Oxford as emmy . network Fellow. James Worrell was supported by EPSRC Fellowship EP/X033813/1.

This work partly arose through exchanges held at the workshop *Algorithmic Aspects of Dynamical Systems*, which took place in April and May 2023 at McGill University's Bellairs Research Institute in Barbados.

REFERENCES

- [1] Boris Adamczewski and Yann Bugeaud. 2007. On the complexity of algebraic numbers I. Expansions in integer bases. *Annals of Mathematics* (2007), 547–565.
- [2] Jean-Paul Allouche and Jeffrey Shallit. 2003. *Automatic sequences: theory, applications, generalizations*. Cambridge university press.
- [3] Pierre Arnoux, Christian Mauduit, Iekata Shiokawa, and Jun-ichi Tamura. 1994. Complexity of sequences defined by billiard in the cube. *Bulletin de la Société Mathématique de France* 122, 1 (1994), 1–12.
- [4] Yu. Baryshnikov. 1995. Complexity of trajectories in rectangular billiards. *Communications in mathematical physics* 174 (1995), 43–56.
- [5] Paul T Bateman, Carl G Jockusch, and Alan R Woods. 1993. Decidability and undecidability of theories with a predicate for the primes. *The Journal of symbolic logic* 58, 2 (1993), 672–687.
- [6] Nicolas Bedaride. 2009. Directional complexity of the hypercubic billiard. *Discrete mathematics* 309, 8 (2009), 2053–2066.
- [7] Valérie Berthé, Toghrol Karimov, Joël Ouaknine, Mihir Vahanwala, and James Worrell. 2023. The Monadic Theory of Toric Words. *arXiv preprint arXiv:2311.04895* (2023).
- [8] Achim Blumensath. 2023. Monadic Second-Order Model Theory. www.fi.muni.cz/~blumens/. [Online; accessed on 09 November 2023].
- [9] J. R. Büchi. 1990. On a Decision Method in Restricted Second Order Arithmetic. In *The Collected Works of J. Richard Büchi*. Springer New York, 425–435.
- [10] J Richard Büchi and Lawrence H Landweber. 1969. Definability in the monadic second-order theory of successor1. *The Journal of Symbolic Logic* 34, 2 (1969), 166–170.
- [11] Jin-yi Cai, Richard J. Lipton, and Yechezkel Zalcstein. 2000. The Complexity of the A B C Problem. *SIAM J. Comput.* 29, 6 (2000), 1878–1888. [arXiv:https://doi.org/10.1137/S0097539794276853](https://doi.org/10.1137/S0097539794276853)
- [12] Olivier Carton and Wolfgang Thomas. 2002. The Monadic Theory of Morphic Infinite Words and Generalizations. *Information and Computation* 176, 1 (2002), 51–65.
- [13] Henri Cohen. 2013. *A course in computational algebraic number theory*. Vol. 138. Springer Science & Business Media.
- [14] George B Dantzig, B Curtis Eaves, et al. 1973. Fourier-Motzkin elimination and its dual. *J. Comb. Theory, Ser. A* 14, 3 (1973), 288–297.
- [15] Calvin C. Elgot and Michael O. Rabin. 1966. Decidability and Undecidability of Extensions of Second (first) Order Theory of (generalized) Successor. *The Journal of Symbolic Logic* 31, 02 (1966), 169–181.
- [16] Graham Everest, Alf van der Poorten, Igor Shparlinski, and Thomas Ward. 2003. *Recurrence Sequences*. American Mathematical Society.
- [17] N Pytheas Fogg, Valérie Berthé, Sébastien Ferenczi, Christian Mauduit, and Anne Siegel. 2002. *Substitutions in dynamics, arithmetics and combinatorics*. Springer.
- [18] Steven M. Gonek and Hugh L. Montgomery. 2016. Kronecker's Approximation Theorem. *Indagationes Mathematicae* 27, 2 (2016), 506–523. <https://www.sciencedirect.com/science/article/pii/S0019357716000148> In Memoriam J.G. Van der Corput (1890–1975) Part 2.
- [19] Glyn Harman. 2002. One hundred years of normal numbers. In *Surveys in Number Theory*. AK Peters/CRC Press, 57–74.
- [20] Toghrol Karimov, Engel Lefauchaux, Joël Ouaknine, David Purser, Anton Varonka, Markus A. Whiteland, and James Worrell. 2022. What's decidable about linear loops? *Proc. ACM Program. Lang.* 6, POPL (2022), 1–25.
- [21] Serge Lang. 1966. *Introduction to Transcendental Numbers*. Addison-Wesley.
- [22] M. Lothaire. 2002. *Algebraic combinatorics on words*. Vol. 90. Cambridge university press.
- [23] A. Macintyre and A. J. Wilkie. 1996. On the Decidability of the Real Exponential Field. In *Kreisliana. About and Around Georg Kreisel*, Piergiorgio Odifreddi (Ed.). AK Peters, 441–467.
- [24] David W Masser. 1988. Linear relations on algebraic groups. *New Advances in Transcendence Theory* (1988), 248–262.
- [25] Eugene M Matveev. 2000. An explicit lower bound for a homogeneous rational linear form in the logarithms of algebraic numbers. II. *Izvestiya: Mathematics* 64, 6 (2000), 1217.
- [26] An. Muchnik, A. Semenov, and M. Ushakov. 2003. Almost periodic sequences. *Theoretical Computer Science* 304, 1 (2003), 1–33.
- [27] Martine Queffelec. 2006. Old and new results on normality. *Lecture Notes-Monograph Series* (2006), 225–236.
- [28] Alexander Rabinovich. 2007. On decidability of monadic logic of order over the naturals extended by monadic predicates. *Information and Computation* 205, 6 (2007), 870–889.
- [29] Alexander Rabinovich and Wolfgang Thomas. 2006. Decidable theories of the ordering of natural numbers with unary predicates. In *International Workshop on Computer Science Logic*. Springer, 562–574.
- [30] Andrzej Schinzel and Robert Tijdeman. 1976. On the equation $y^m = P(x)$. *Acta Arithmetica* 31 (1976), 199–204.
- [31] Aleksei L'vovich Semenov. 1980. On certain extensions of the arithmetic of addition of natural numbers. *Mathematics of The USSR-Izvestiya* 15 (1980), 401–418. <https://api.semanticscholar.org/CorpusID:123422158>
- [32] A. L. Semënov. 1984. Logical Theories of One-Place Functions on the Set of Natural Numbers. *Mathematics of the USSR-Izvestiya* 22, 3 (1984), 587–618.
- [33] Wolfgang Thomas. 1997. Languages, automata, and logic. In *Handbook of Formal Languages: Volume 3 Beyond Words*. Springer, 389–455.
- [34] Michel Waldschmidt. 2000. *Diophantine Approximation on Linear Algebraic Groups: Transcendence Properties of the Exponential Function in Several Variables*. Springer Berlin Heidelberg, Berlin, Heidelberg.

A OMITTED PROOFS

A.1 Proof of Lemma 2.9

We will use the following lemmas to prove Lem. 2.9.

LEMMA A.1. *Assume that $d, e \geq 1$ and that*

$$f(x_1, \dots, x_e) = \frac{c_1}{\sum_{i=1}^e b_{i,1}x_i} + \dots + \frac{c_d}{\sum_{i=1}^e b_{i,d}x_i}$$

for some $c_j, b_{i,j} \in \mathbb{Q}$ such that for $1 \leq j < j' \leq d$, $\sum_{i=1}^e b_{i,j}x_i \neq s \sum_{i=1}^e b_{i,j'}x_i$ does not hold for any $s \in \mathbb{R}$. If $f(x_1, \dots, x_d) = 0$, then $c_1, \dots, c_d = 0$.

PROOF. For $1 \leq j \leq d$, Let V_j be the $e-1$ -dimensional subspace of \mathbb{R}^e where $\sum_{i=1}^e b_{i,j}x_i = 0$ holds for all $(x_1, \dots, x_e) \in \mathbb{R}^e$. Assume that for $1 \leq j \leq d$, $c_j \neq 0$.

If $(\mathbf{x}_n)_{n=0}^\infty$ is a \mathbb{R}^e -valued sequence converging to $\mathbf{x} \in V_j$, then $\lim_{n \rightarrow \infty} |f(\mathbf{x}_n)| = +\infty$. Hence, as $f(x_1, \dots, x_e) = 0$, $\mathbf{x} \in V_{j'}$ for some $1 \leq j' \leq d$ unequal to j . Then $V_j \cap V_{j'}$ is again a linear subspace of \mathbb{R}^e of dimension at most $e-1$.

If $V_j \cap V_{j'}$ has dimension less than $e-1$ for all $1 \leq j' \leq d$ unequal to j , then

$$V_j \subsetneq \bigcup_{\substack{1 \leq j' \leq d \\ j' \neq j}} (V_j \cap V_{j'})$$

giving a contradiction that each $\mathbf{x} \in V_j$ is in some $V_{j'}$ with $j' \neq j$.

Thus, $V_j = V_{j'}$ for some $j' \neq j$, and so $\sum_{i=1}^e b_{i,j}x_i = 0$ if and only if $\sum_{i=1}^e b_{i,j'}x_i = 0$. Hence, there is some $s \in \mathbb{R}_{\neq 0}$ such that $b_{i,j} = sb_{i,j'}$ for all $1 \leq i \leq e$. This gives a contradiction, and so $c_j = 0$. As this holds for all $1 \leq j \leq d$, the lemma follows. \square

LEMMA A.2. *Assume that $\lambda_1, \dots, \lambda_d \in \mathbb{R}_{>1} \cap \overline{\mathbb{Q}}$ are pairwise multiplicatively independent. Then $1/\log(\lambda_1), \dots, 1/\log(\lambda_d)$ are linearly independent over \mathbb{Q} .*

PROOF. First, using Th. 2.4, compute a basis for $G_M(\lambda_1, \dots, \lambda_d)$ and select a maximum multiplicative independent subset $\{\lambda_1, \dots, \lambda_e\}$, possibly needing to renumber the λ_i . Then, for all $1 \leq j \leq d$, one can compute $b_{i,j}$ for $1 \leq i \leq e$ such that $\log(\lambda_j) = \sum_{i=1}^e b_{i,j} \log(\lambda_i)$ using the known multiplicative relationships. Then we have to show that when $c_1, \dots, c_d \in \mathbb{Q}$ and

$$\frac{c_1}{\sum_{i=1}^e b_{i,1} \log(\lambda_i)} + \dots + \frac{c_d}{\sum_{i=1}^e b_{i,d} \log(\lambda_i)} = 0 \quad (12)$$

Schanuel's conjecture implies that all c_i are zero. By multiplying (12) by $\prod_{j=1}^d \sum_{i=1}^e b_{i,j} \log(\lambda_i)$, we obtain a polynomial expression in $\log(\lambda_1), \dots, \log(\lambda_e)$ that equals zero.

We apply Schanuel's conjecture (Conj. 2.7) with $\alpha_i = \log(\lambda_i)$ for $1 \leq i \leq e$. As $\lambda_1, \dots, \lambda_e$ are multiplicatively independent, $\log(\lambda_1), \dots, \log(\lambda_e)$ are linearly independent over \mathbb{Q} by Baker's theorem (Thm. 2.10). Then $\{\log(\lambda_1), \dots, \log(\lambda_e), \lambda_1, \dots, \lambda_e\}$ has transcendence degree at least d by Schanuel's conjecture while also being equal to the transcendence degree of $\{\log(\lambda_1), \dots, \log(\lambda_e)\}$ as all λ_i are algebraic. Hence, $\log(\lambda_1), \dots, \log(\lambda_e)$ are algebraically independent.

This implies that the polynomial expression obtained from (12), has to evaluate trivially to zero. That is, the rational function

$$f(x_1, \dots, x_e) = \frac{c_1}{\sum_{i=1}^e b_{i,1}x_i} + \dots + \frac{c_d}{\sum_{i=1}^e b_{i,d}x_i}$$

is exactly zero. Assume that $1 \leq j < j' < d$ and $\sum_{i=1}^e b_{i,j}x_i \neq s \sum_{i=1}^e b_{i,j'}x_i$ holds for a real number s . As $\lambda_{j'} \neq 0$, some $b_{i,j'}$ is non-zero. Then $s = b_{i,j}/b_{i,j'} \in \mathbb{Q}$. Hence,

$$\log(\lambda_j) = \sum_{i=1}^e b_{i,j} \log(\lambda_i) = \sum_{i=1}^e sb_{i,j'} \log(\lambda_i) = s \log(\lambda_{j'}),$$

contradicting that λ_j and $\lambda_{j'}$ are multiplicatively independent. Thus, the hypothesis of Lem. A.1 is satisfied and all c_j are 0. We conclude the statement. \square

Now Lem. 2.9 easily follows.

PROOF OF LEM. 2.9. If λ_i and λ_j are multiplicatively dependent, say $\lambda_i^a = \lambda_j^b$ for some non-zero integers a and b , then $a \log(\lambda_i) = b \log(\lambda_j)$. Hence, $a/\log(\lambda_j) = b/\log(\lambda_i)$, giving a non-trivial element in $G_A(1/\log(\lambda_1), \dots, 1/\log(\lambda_d))$.

Meanwhile, by Lem. A.2, for any pairwise multiplicative independent subset of $\{\lambda_1, \dots, \lambda_d\}$, the reciprocals of their logs are linearly independent.

Together, this implies that any \mathbb{Q} -linear relationship among $1/\log(\lambda_1), \dots, 1/\log(\lambda_d)$ can be reduced to a relationship generated by pairwise multiplicative relationships. One can easily find a basis among these by computing for each pair (λ_i, λ_j) whether they multiplicatively dependent using Thm. 2.4. \square

A.2 Proof of Lemma 2.11

PROOF OF LEM. 2.11. By the two assumptions, for any $1 \leq i < j < k \leq d$ there exist $b_1, b_2, b_3 \in \mathbb{Z}_{\neq 0}$ such that $\lambda_i^{b_1} \lambda_j^{b_2} \lambda_k^{b_3} = 1$. Equivalently, $b_1 \log(\lambda_i) + b_2 \log(\lambda_j) + b_3 \log(\lambda_k) = 0$. Hence, for $1 \leq j \leq d$, let $b_{1,j}, b_{2,j} \in \mathbb{Q}$ be such that $\log(\lambda_j) = b_{1,j} \log(\lambda_1) + b_{2,j} \log(\lambda_2)$. Then $b_{1,1} = b_{2,2} = 0$ and all other $b_{i,j}$ are non-zero.

For a contradiction, let $c_1, \dots, c_d \in \mathbb{Q}$ be rational numbers such that $\sum_{j=1}^d c_j / \log(\lambda_j) = 0$. Multiplying by $\prod_{j=1}^d \log(\lambda_j)$ gives

$$\sum_{j=1}^d c_j \prod_{i=1, i \neq j}^d (b_{1,i} \log(\lambda_1) + b_{2,i} \log(\lambda_2)) = 0,$$

which simplifies to

$$\sum_{i=0}^d e_i \log(\lambda_1)^i \log(\lambda_2)^{d-i} = 0 \quad (13)$$

for some $e_i \in \mathbb{Q}$. Assume not all e_i are zero. Then dividing by $\log(\lambda_2)^d$ shows that $\log(\lambda_1)/\log(\lambda_2)$ is the root of the non-zero polynomial $\sum_{i=0}^d e_i x^i \in \mathbb{Q}[x]$. That is, $\log(\lambda_1)/\log(\lambda_2)$ is an algebraic number, say α and so $\log(\lambda_1) - \alpha \log(\lambda_2) = 0$, contradicting Baker's theorem (Thm. 2.10). Hence, all e_i have to be zero.

As (13) is obtained by multiplying $\sum_{j=1}^d \frac{c_j}{\log(\lambda_j)}$ with the non-zero number $\prod_{j=1}^d \log(\lambda_j)$,

$$\sum_{j=1}^d \frac{c_j}{b_{1,j}x_1 + b_{2,j}x_2} = 0 \quad (14)$$

As in Lem. A.2, we deduce that as all λ_i are pairwise multiplicatively dependent and so (14) satisfies the hypothesis of Lem. A.1. Thus, all c_i are zero and the statement follows. \square

A.3 Proof of Theorem 2.13

PROOF OF THM. 2.13. If ρ_1 and ρ_2 are multiplicatively dependent, say $\rho^{m_1} = \rho_2^{m_2}$, let $\rho_3 = \rho_1^{1/m_2}$, $R_3 = \max(R_1^{1/m_2}, R_2^{1/m_1})$. Then $R_3 < \rho_3$ and solving

$$|c_1 \rho_3^{n_1'} - c_2 \rho_3^{n_2'}| \leq (b_1 + b_2) R_3^{\max(n_1 + n_2)}$$

when $c_1 \rho_3^{n_1'} \neq c_2 \rho_3^{n_2'}$ gives all solutions to (3) by setting $n_1 = m_2 n_1'$ and $n_2 = m_1 n_2'$. As the left-hand side can be bounded from below by $C_1 \rho_3^{\max(n_1', n_2')}$ for a computable constant C_1 when it is non-zero, $\max(n_1', n_2')$ can be bounded from above and so n_1 and n_2 as well.

If ρ_1 and ρ_2 are multiplicatively independent, it is sufficient to bound n_1, n_2 that satisfy

$$|c_1 \rho_1^{n_1} - c_2 \rho_2^{n_2}| \geq 2b_1 R_1^{n_1}, 2b_2 R_2^{n_2} \quad (15)$$

as adding the two cases gives the result. We can assume that $\rho_1 < \rho_2$. After dividing both sides by $c_1 \rho_1^{n_1}$, we apply Matveev's result to $\Lambda = c_1^{-1} c_2 (\rho_1^{-1})^{n_1} \rho_2^{n_2} - 1$ to find that

$$\log |\Lambda| \geq -C_1 (1 + \log(\max(1, n_1, n_2)))$$

for some computable constant $C_1 > 0$. Thus, we want n_1 and n_2 to satisfy

$$\begin{aligned} \log |\Lambda| &\geq \log(2b_1/c_1) + n_1 \log(\rho_1/R_1) \quad \text{and} \\ \log |\Lambda| &\geq \log(2b_2/c_1) + n_2 \log(\rho_2) - n_1 \log(R_1). \end{aligned} \quad (16)$$

If $n_1 \geq n_2$ and the first of these equations is false, then

$$C_1 (1 + \log(\max(1, n_1))) \geq \log(c_2/(2b_1)) + n_1 \log(\rho_1/R_1) \quad (17)$$

We can effectively bound n_1 (and thus n_2) as the right-hand side grows logarithmically while the left-hand grows linearly in n_1 .

If $n_2 \geq n_1$ and the second equation of (16) is false,

$$C_1 (1 + \log(\max(1, n_2))) \leq \log(c_2/(2b_1)) + n_1 \log(\rho_1/R_1).$$

Thus, we can compute constants C_3, C_4 such that $C_4 > 0$ and if the second equation of (16) is false, then $n_1 \geq C_3 + C_4 \log(n_2)$. Thus, we can assume that $n_1 \leq C_3 + C_4 \log(n_2)$. If the second equation of (16) is false,

$$C_1 (1 + \log(\max(1, n_2))) \geq \log(c_1/2b_2) + n_2 \log(\rho_2) - n_1 \log(R_1).$$

Using our bound on n_1 , we obtain that when $n_2 \geq 1$

$$C_1 (1 + \log(n_2)) \geq \log(c_1/2b_2) + n_2 \log(\rho_2) - (C_3 + C_4 \log(n_2)) \log(R_1).$$

The left-hand side grows logarithmically in n_2 while the right-hand side grows linearly. Hence, we can again bound n_2 (and thus n_1). The result follows. \square

A.4 Proof of Theorem 4.18

PROOF OF THM. 4.18. Consider a given deterministic Muller automaton \mathcal{A} as a directed graph allowing multiple edges. We partition the graph into its strongly connected components (SCCs) and call an SCC without outgoing edges a bottom SCC. We will show that the set of states visited infinitely often by the run of \mathcal{A} on α is precisely a bottom SCC. Hence, we decide Acc_α by simulating this run until a bottom SCC is inevitably reached. Then α is accepted if and only if this bottom SCC is in the Muller acceptance condition.

We need to show that (a) if an SCC is not a bottom SCC, then the run eventually exits it; and (b) if the run enters a bottom SCC, it visits all its states infinitely often.

For (a), let S be a non-bottom SCC. There thus exist $q_1 \in S, b \in \Sigma$ such that $\delta(q_1, b) \notin S$, i.e. reading the letter b in state q_1 exits S . We will order the states of S as q_1, \dots, q_k , construct words $u_1 = b, \dots, u_k \in \Sigma^+$, and inductively prove that for all $j \leq i, \delta(q_j, u_i) \notin S$. Since α is weakly normal, u_k will inevitably occur as a factor, and hence, the run will exit S .

We have observed the base case to hold with $u_1 = b$. For the induction step, assume that for all $j \leq i, \delta(q_j, u_i) \notin S$. Now, if $\delta(q_{i+1}, u_i) \notin S$, take $u_{i+1} = u_i$. Else, if $\delta(q_{i+1}, u_i) = q \in S$, the strong connectivity of S implies that $\delta(q, v_i) = q_1$ for some $v_i \in \Sigma^*$. Thus, take $u_{i+1} = u_i v_i b$, and observe that for all $j \leq i+1, \delta(q_j, u_{i+1}) \notin S$.

We prove (b) similarly. Fix an order of states q_1, \dots, q_k of S . By definition, a run entering the bottom SCC S will be confined in S . It thus suffices to prove that for any $q \in S$, we can inductively construct a word $u_k \in \Sigma^+$ such that for all $j \leq k$, the non-empty run of u_k on \mathcal{A} starting from q_j visits q . The induction is similar to the one above. Choose $q_1 \in S, u_1 \in \Sigma^*$ to be such that $\delta(q_1, u_1) = q$. By the induction hypothesis, for every $j \leq i$, the run on u_i starting in q_j visits q . If the run on u_i starting in q_{i+1} visits q , take $u_{i+1} = u_i$. Else, use that S is a bottom SCC to identify $v_i \in \Sigma^+$ such that $\delta(q_{i+1}, u_i v_i) = q$, and take $u_{i+1} = u_i v_i$. We have thus ensured that for all $j \leq i+1$, the run on u_i starting in q_j visits q . Since α is weakly normal, u_k occurs as a factor infinitely often, and hence all $q \in S$ are visited infinitely often. \square

A.5 Proof of Lemma 5.4

PROOF OF LEM. 5.4. We first deal with the first part of (d). Fix distinct i, j . Recall we assumed there were only finitely many pairs $(n_i, n_j) \in \mathbb{N}^2$ such that $c_i \rho_i^{n_i} = c_j \rho_j^{n_j}$. Using Thm. 2.4, we can compute a basis for $G_M(c_i/c_j, \rho_i, \rho_j)$ and with linear algebra, we can compute all such (k_1, k_2, k_3) such that $k_1 = 1$. These elements are exactly the solutions of $c_i \rho_i^{k_2} = c_j \rho_j^{-k_3}$ in integers. As there are only finitely many such k_2 and k_3 , we can compute and set $k_{i,j} = \max(k_2, -k_3)$.

We continue with the proof of (a-e). We already have a bound for the first part of (d). Let us treat (e). For $1 \leq i \leq d$, apply Lem. 2.5 to find $b_i, R_i > 0$ such that $|u_n^{(i)} - c_i \rho_i^n| < b_i R_i^n$ for all $n \geq 0$. Then,

$$\begin{aligned} u_{n_i}^{(i)} - u_{n_j}^{(j)} &= c_i \rho_i^n - c_j \rho_j^n + u_{n_i}^{(i)} - c_i \rho_i^n - u_{n_j}^{(j)} + c_j \rho_j^n \\ &> c_i \rho_i^n - c_j \rho_j^n - b_j R_j^n - b_j R_j^n. \end{aligned}$$

Using Thm. 2.13, for each pair $1 \leq i < j \leq d$, we compute $k'_{i,j}$ such that for all $n_i, n_j \geq k_{i,j}$, $|c_i \rho_i^n - c_j \rho_j^n| > b_i R_i^n + b_j R_j^n$ such that for such $n_i, n_j, u_{n_i}^{(i)} - u_{n_j}^{(j)}$ and $c_i \rho_i^n - c_j \rho_j^n$ have the same sign. Thus, (e) follows.

For the second part of (d), assume that $n_i, n_j \geq k_{i,j}, k'_{i,j}$. Now note the first part of (d) implies that, $c_i \rho_i^{n_i} > c_j \rho_j^{n_j}$ or $c_i \rho_i^{n_i} < c_j \rho_j^{n_j}$ and thus by (e), $u_{n_i}^{(i)} \neq u_{n_j}^{(j)}$.

For (a), let $1 \leq i \leq d$. Then,

$$\begin{aligned} u_{n+1}^{(i)} - u_n^{(i)} &< c_i \rho_i^{n+1} - c_i \rho_i^{n+1} - b_i R_i^{n+1} - b_i R_i^n \\ &= c_i (\rho_i - 1) \rho_i^n - b_i (R_i + 1) R_i^n. \end{aligned}$$

As $\rho_i > R_i$, one can compute a bound k_i such that $c_i (\rho_i - 1) \rho_i^n > b_i (R_i + 1) R_i^n$ for all $n \geq k_i$.

Thus, when $m_i \geq k_i, k_{i,j}, k'_{i,j}$ for all $1 \leq i, d \leq d$, (a), (d) and (e) are satisfied. Let m'_i satisfy these conditions. Then choose a $N' \geq 0$ such that $N' \geq u_{m'_i}^{(i)}$ for all $1 \leq i \leq d$. Then let m_1 be the smallest number such that $u_{m_1}^{(1)} \geq N'$ and $N = u_{m_1}^{(1)}$. Then for $1 \leq i \leq d$, let m_i be the smallest number such that $u_{m_i}^{(i)} \geq N$. Then (b) and (c) are satisfied. \square

A.6 Proof of Lemma 6.4

PROOF OF LEM. 6.4. Recall that every finite factor of a uniformly recurrent word occurs infinitely often and with bounded gaps. Let w be a finite pattern that occurs in α . We will show how to compute $M \in \mathbb{N}$ such that w occurs at least twice in every factor of α of length M . The value M is then an upper bound on the gaps between consecutive occurrences of w in α . Note that such M exists by the uniform recurrence assumption. Let T_n be the set of all factors of α of length n , which can be computed by enumerating all words of length n and checking whether each one occurs in α . The value M can be found by computing T_n for increasing values of n . \square

A.7 Proof of Lemma 6.5

PROOF OF LEM. 6.5. For $b = 1b(1) \cdots b(m) \in \Sigma$, let

$$S_{b,k} = \{x \in \mathbb{T}^{d-1} : g^{(k)}(x) \in S_b\}.$$

Since $g: \mathbb{T}^{d-1} \rightarrow \mathbb{T}^{d-1}$ is a homeomorphism, $S_{b,k}$ is an open subset of \mathbb{T}^{d-1} . Since $g^{(k)}(x_2, \dots, x_d) = (\{x_2 + k\delta_2\}, \dots, \{x_d + k\delta_d\})$, from (10) it follows that $S_{b,k}$ is the set of all $(x_2, \dots, x_d) \in \mathbb{T}^{d-1}$ satisfying

$$\frac{\{x_{b(m)} + k\delta_{b(m)}\}}{\delta_{b(m)}} < \dots < \frac{\{x_{b(1)} + k\delta_{b(1)}\}}{\delta_{b(1)}} < 1$$

and

$$\bigwedge_{\substack{1 \leq j \leq d \\ j \neq b(1), \dots, b(m)}} \frac{\{x_j + k\delta_j\}}{\delta_j} > 1.$$

For $2 \leq i \leq d$, let $t_i = \lfloor k\delta_i \rfloor$. Observe that $\{x_i + k\delta_i\} = x_i + k\delta_i - t_i$ if $x_i + k\delta_i < t_i + 1$ and $\{x_i + k\delta_i\} = x_i + k\delta_i - (t_i + 1)$ otherwise. Moreover, $x_i + k\delta_i < t_i + 1$ is equivalent to $x_i/\delta_i + k < \frac{t_i + 1}{\delta_i}$. Therefore,

$\frac{\{x_j + k\delta_j\}}{\delta_j} \succ 1$ (where \succ is an (in)equality symbol) is equivalent to

$$\frac{x_i}{\delta_i} + k < \frac{t_i + 1}{\delta_i} \Rightarrow \frac{x_j + k\delta_j - t_j}{\delta_j} \succ 1 \quad \wedge$$

$$\frac{x_i}{\delta_i} + k \geq \frac{t_i + 1}{\delta_i} \Rightarrow \frac{x_j + k\delta_j - (t_j + 1)}{\delta_j} \succ 1.$$

Rearranging, this formula can be written as a Boolean combination of inequalities of the form (11) where h is a \mathbb{Q} -affine form. Similarly, $\frac{\{x_i + k\delta_i\}}{\delta_i} < \frac{\{x_j + k\delta_j\}}{\delta_j}$ can be equivalently written as a Boolean combination of inequalities of the form (11) by conditioning on whether $x_i/\delta_i + k < (t_i + 1)/\delta_i$ and $x_j/\delta_j + k < (t_j + 1)/\delta_j$. Finally, observe that $0 \leq x_i < 1$ is equivalent to $0 \leq x_i\delta_i < 1/\delta_i$. We conclude that $S_{b,k}$ can be defined by a Boolean combination of inequalities of the form (11).

It remains to define $S_w = \bigcap_{k=0}^{|w|-1} S_{w(k),k}$. Since each $S_{w(k),k}$ is open and defined by a Boolean combination of inequalities of the form (11), we conclude the same for S_w . \square

A.8 Proof of Theorem 7.1

PROOF OF COR. 7.1. By assumption, $p/q = A^d/B^d$ for some coprime integers $A, B \geq 1$. Thus, $p = qA^d/B^d$ and B^d divides q . Let $f_1(n) = (q/B^d)n^d$ and $f_2(n) = Ab^n$. Then $g_1(n) = f_1(n) = (q/B^d)n^d$ and $g_2(n) = g_1 \circ f_2(n) = g_1(Ab^n) = (q/B^d)A^d b^{nd} = pb^{nd}$. Let β be the characteristic word of $(\{g_1(n) : n \in \mathbb{N}\}, \{g_2(n) : n \in \mathbb{N}\})$. Then the problem Acc_β is decidable by Thm. 4.11 as by [12, Theorem 5.2], $f_1, f_2 \in \mathcal{F}$.

Then we construct a transducer \mathcal{B} from β to α as follows. We keep track of the number of occurrences of $(1, 0)$ and $(1, 1)$ we have encountered modulo B^d . When this counter is zero modulo B^d , we output $\beta(n)$, otherwise we output $(0, j)$ instead of (i, j) . Thus, the second predicate remains unchanged while for the first we only count every B^d th term. Hence, $\alpha = \mathcal{B}(\beta)$ and so Acc_α is decidable using Lem. 4.5. \square

A.9 Proof of Theorem 7.4

PROOF OF THM. 7.4. The proof mirrors the proof of Thm. 7.2 closely.

To show that Acc_β reduces to Acc_α , let α' be the word obtained from n^2 and $b \cdot b^{2n}$. Then the transducer \mathcal{B} that changes every $(1, 1)$ into $(1, 0)$ (and leaves everything else unchanged) has the property that $\mathcal{B}(\alpha) = \alpha'$. This follows from the fact that a power of b is not a square if and only if it is of the form $b \cdot b^{2n}$. By Lem. 4.5, $\text{Acc}_{\alpha'}$ reduces to Acc_α . As $\text{Acc}_{\alpha'}$ and Acc_β are Turing-equivalent, one direction follows.

To show the other direction, let $\gamma \in \{(0, 1), (1, 0), (1, 1)\}^\omega$ be the word obtained by deleting all occurrences of $(0, 0)$ from α . Denoting $\mathbf{0} = (0, 0)$, we have:

$$\alpha = \mathbf{0}^{k_0} \gamma(0) \cdots \mathbf{0}^{k_n} \gamma(n) \cdots$$

We invoke Lem. 7.3 to argue that for any K , we can compute N such that for all $n \geq N$, $k_n > K$. The number of intervening $\mathbf{0}$'s is effectively eventually lower bounded, and thanks to the procyclic nature of the predicates, can still be computed modulo any m . Thus, similar to the proof of Thm. 4.15, one can use Cor. 4.7 to show that Acc_α reduces to Acc_γ . Our strategy to reduce Acc_γ to Acc_β is to closely follow the proof of Thm. 7.2 (3), and apply Cor. 4.7 in a similar way.

The terms $(1, 0)$ signify the squares which are not powers of b , $(0, 1)$, the numbers $b \cdot b^{2n}$ and $(1, 1)$ the numbers b^{2n} . We express γ as

$$\gamma = (1, 0)^{k_0} (1, 1) (1, 0)^{k_1} (0, 1) \cdots (1, 0)^{k_{2n}} (1, 1) (1, 0)^{k_{2n+1}} (0, 1) \cdots$$

In order to apply Cor. 4.7, we are interested in computing k_n . Similar to the proof of Thm. 7.2 (3), we record:

$$\begin{aligned}
 S_{2n} &= n + \sum_{i=0}^{2n} k_i = b^n \\
 S_{2n+1} &= n + \sum_{i=0}^{2n+1} k_i = \lfloor b^n \sqrt{b} \rfloor \\
 k_{2n} &= b^n - \lfloor b^{n-1} \sqrt{b} \rfloor - 1 \\
 k_{2n+1} &= \lfloor b^n \sqrt{b} \rfloor - b^n \\
 \beta(n) &\equiv \lfloor b^n \sqrt{b} \rfloor \pmod{b} \\
 S_{2n+2} &= bS_{2n} = S_{2n+1} + k_{2n+1} + 1 \\
 S_{2n+3} &= bS_{2n+1} + \beta(n+1) = S_{2n+2} + k_{2n+3}
 \end{aligned}$$

Using these, one can keep track of k_n modulo any m given the digits of β . The rest of the proof, which goes on to appropriately invoke Cor. 4.7, proceeds similarly, and we omit it. \square